

**URINARY S/O**

**URINARY S/O**  
MODERATE CALORIE

**URINARY S/O**  
HYPOALLERGENIC

**URINARY S/O**  
BLADDER COMFORT

**URINARY S/O**  
AGEING<sup>7+</sup>  
BLADDER COMFORT

**URINARY S/O**  
BLADDER COMFORT  
SATIETY



**URINARY S/O**

**URINARY S/O**  
SMALL DOGS  
S

**URINARY U/C**

**URINARY S/O**  
MODERATE CALORIE

**URINARY S/O** AGEING<sup>7+</sup>

**URINARY S/O**  
SATIETY

**URINARY S/O**  
HYPOALLERGENIC


This resource has been produced with the greatest care and incorporates the most up-to-date scientific information and research findings. We recommend that you refer to our packaging as the primary reference for product information as this can change with time. Bearing in mind the diversity and complexity of clinical cases in dogs and cats, it is important to note that the suggestions in this resource for further tests and therapeutic options cannot be exhaustive. Under no circumstances should the treatments and dietary solutions suggested here replace examination by a veterinary surgeon.

The publishers and the authors will accept no responsibility for the failure of any of the treatments or nutritional solutions put forward in this resource. Not for use in Canada and in USA.

While the information is considered to be true and correct at the date of publication (**May 25**), changes in circumstances after the time of publication may impact on the accuracy of the information.



# FELINE NUTRITIONAL MANAGEMENT DECISION TREE





URINARY S/O

# URINARY S/O



Available in  
1.5, 3.5, 7  
and 9kg bags

## RECOMMENDED DAILY INTAKE & FEEDING DURATION

ADULT CAT	CAT'S WEIGHT (kg)	BODY CONDITION		
		THIN	NORMAL	OVERWEIGHT
		grams	grams	grams
	2	39	33	26
	2.5	46	38	31
	3	52	44	35
	3.5	59	49	39
	4	64	54	43
	4.5	70	58	47
	5	75	63	50
	5.5	81	67	54
	6	86	72	57
	6.5	91	76	61
	7	96	80	64
	7.5	101	84	67
	8	105	88	70
	8.5	110	92	73
	9	115	96	76
	9.5	119	99	79
	10	123	103	82

Urinary S/O is a complete dietetic feed for cats, formulated to dissolve struvite stones and reduce their recurrence through its urine acidifying properties and its low level of magnesium.

**RECOMMENDATIONS:** It is recommended that a veterinarian's opinion be sought before use. Feed Urinary S/O for 5 to 12 weeks for the dissolution of struvite stones and up to 6 months for the reduction of struvite stone recurrence. The cat should be re-evaluated regularly by the veterinarian to assess the relevance of a lifelong recommendation. Water should be available at all times.

**NOTE:** The wet version of Feline Urinary S/O is recommended for recurrent idiopathic cystitis. In older cats, renal function should be checked before recommending a Urinary S/O diet. The use of Urinary Ageing 7+ Bladder Comfort should be considered in older cats.

## LOWER URINARY TRACT DISEASE

### RECOMMENDED FOR CASES OF

- Struvite uroliths: dissolution and management of recurrence
- Calcium oxalate uroliths: management of recurrence

### NOT RECOMMENDED IN CASE OF

- Chronic Kidney Disease (CKD)
- Heart disease (when sodium restriction is sought)
- Concurrent use of urine-acidifying medication
- Growth, gestation/lactation



### STRUVITE DISSOLUTION

Helps dissolve all types of struvite stones.



### URINE DILUTION

Urine dilution makes the urine less liable to form struvite and calcium oxalate stones.



### LOW RSS

Helps lower the concentration of ions contributing to crystal formation.



### LOW MAGNESIUM

Reduced level of magnesium, a natural component of struvite crystals.



This diet is formulated to promote a healthy urinary environment.

### KEY VALUES

	Per 1000 kcal	Per 100g as fed
Moisture	14.20 g	5.50 g
Protein	89.10 g	34.50 g
Fat Content	38.74 g	15.00 g
Crude Fibre	7.49 g	2.90 g
Ash	22.99 g	8.90 g
Carbohydrate (NFE)	85.74 g	33.20 g
Dietary Fibre	17.56 g	6.80 g
Omega-6	8.76 g	3.39 g
Omega-3	2.07 g	0.80 g
EPA + DHA	0.96 g	0.37 g
Calcium	2.32 g	0.90 g
Phosphorus	2.32 g	0.90 g
Sodium	3.36 g	1.30 g
Magnesium	0.13 g	0.05 g
Potassium	2.58 g	1.00 g
Copper	3.87 mg	1.50 mg
Chloride	5.17 g	2.00 g
Vitamin A	6198 IU	2400 IU
Vitamin D	207 IU	80 IU

Metabolisable energy\* 387.20 Kcal

\*Calculated according to NRC 2006 CF.

### COMPOSITION

Rice, wheat gluten\*\*, dehydrated poultry protein, maize flour, animal fats, hydrolysed animal proteins, maize gluten, minerals, vegetable fibres, fish oil, soya oil, fructo-oligosaccharides, marigold meal.

\*\*L.I.P.: protein selected for its very high digestibility

### ADDITIVES (KG)\*\*

Nutritional additives: Vitamin A: 21500 IU, Vitamin D3: 800 IU, Iron (3b103): 37 mg, Iodine (3b201, 3b202): 3.7 mg, Copper (3b405, 3b406): 11 mg, Manganese (3b502, 3b504): 48 mg, Zinc (3b603, 3b605, 3b606): 136 mg, Selenium (3b801, 3b811, 3b812): 0.06 mg - Antioxidants.

\*\*\* Values reflect only levels added to the formula, not those naturally occurring in components of the diet.



# URINARY S/O



MORSELS IN GRAVY

Available in box  
of 12 pouches  
of 85g

## RECOMMENDED DAILY INTAKE & FEEDING DURATION

ADULT CAT	CAT'S WEIGHT (kg)	BODY CONDITION					
		THIN		NORMAL		OVERWEIGHT	
		grams	pouch	grams	pouch	grams	pouch
	2	170	2	140	1 + 1/2	110	1 + 1/2
	2.5	195	2 + 1/2	165	2	130	1 + 1/2
	3	225	2 + 1/2	185	2	150	2
	3.5	250	3	210	2 + 1/2	165	2
	4	275	3	230	2 + 1/2	185	2
	4.5	300	3 + 1/2	250	3	200	2 + 1/2
	5	320	4	270	3	215	2 + 1/2
	5.5	345	4	285	3 + 1/2	230	2 + 1/2
	6	365	4 + 1/2	305	3 + 1/2	245	3
	6.5	390	4 + 1/2	325	4	260	3
	7	410	5	340	4	275	3
	7.5	430	5	360	4	285	3 + 1/2
	8	450	5 + 1/2	375	4 + 1/2	300	3 + 1/2
	8.5	470	5 + 1/2	390	4 + 1/2	315	3 + 1/2
	9	490	6	410	5	325	4
	9.5	510	6	425	5	340	4
	10	525	6	440	5	350	4

One pouch is equivalent to 18 g of Urinary S/O dry product.

Urinary S/O is a complete dietetic feed for cats, formulated to dissolve struvite stones and reduce their recurrence through its urine acidifying properties.

**RECOMMENDATIONS:** It is recommended that a veterinarian's opinion be sought before use. Feed Urinary S/O for 5 to 12 weeks for the dissolution of struvite stones and up to 6 months for the reduction of struvite stone recurrence. The cat should be re-evaluated regularly by the veterinarian to assess the relevance of a lifelong recommendation. Water should be available at all times.

**NOTE:** In older cats, renal function should be checked before recommending a Urinary S/O diet. The use of Urinary Ageing 7+ Bladder Comfort should be considered in older cats.

## LOWER URINARY TRACT DISEASE

### RECOMMENDED FOR CASES OF

- Struvite uroliths: dissolution and management of recurrence
- Calcium oxalate uroliths: management of recurrence
- Feline idiopathic cystitis (FIC)

### NOT RECOMMENDED IN CASE OF

- Chronic Kidney Disease (CKD)
- Heart disease (when sodium restriction is sought)
- Concurrent use of urine-acidifying medication
- Growth, gestation/lactation



### IDIOPATHIC CYSTITIS

Thanks to the high moisture content of wet food, Urinary S/O Feline supports urine dilution.



### STRUVITE DISSOLUTION

Urinary S/O Feline helps dissolve struvite uroliths.



### LOW RSS

Undersaturated urine creates a urinary environment unfavourable to the development of struvite and calcium oxalate crystals.



### URINE DILUTION

Diluting urine decreases the urinary concentration of struvites and calcium oxalates. The larger urinary volume also supports regular bladder wash-out.



This diet is formulated to promote a healthy urinary environment.

### KEY VALUES

	Per 1000 kcal	Per 100g as fed
Moisture	987 g	80.50 g
Protein	123 g	10 g
Fat Content	42.89 g	3.50 g
Crude Fibre	9.80 g	0.80 g
Ash	25.74 g	2.10 g
Carbohydrate (NFE)	37.99 g	3.10 g
Dietary Fibre	14.71 g	1.20 g
Omega-6	9.80 g	0.80 g
Omega-3	1.96 g	0.16 g
EPA + DHA	1.12 g	0.09 g
Calcium	3.68 g	0.30 g
Phosphorus	2.94 g	0.24 g
Sodium	3.55 g	0.29 g
Magnesium	0.12 g	0.01 g
Potassium	2.21 g	0.18 g
Copper	4.90 mg	0.40 mg
Chloride	5.39 g	0.44 g
Vitamin A	23897 IU	1950 IU
Vitamin D	147 IU	12 IU

Metabolisable energy\* 81.60 Kcal

\*Calculated according to NRC 2006 CF.

### COMPOSITION

Poultry by-products, pork by-products, pork blood products, powder cellulose, corn starch mixture, minerals, vital wheat gluten, maize flour, maize gluten, fish oil, yeasts and part thereof, marigold meal.

### ADDITIVES (KG)\*\*

Nutritional additives: Vitamin D3: 70 IU, Iron (3b103): 7 mg, Iodine (3b202): 0.27 mg, Copper (3b405, 3b406): 2.2 mg, Manganese (3b502, 3b503, 3b504): 2.2 mg, Zinc (3b603, 3b605, 3b606): 22 mg - Technological additives: Sodium bisulphate: 0.5 g.

\*\*\* Values reflect only levels added to the formula, not those naturally occurring in components of the diet.



URINARY S/O



Available in box  
of 12 pouches  
of 85g

#### RECOMMENDED DAILY INTAKE & FEEDING DURATION

	CAT'S WEIGHT (kg)	BODY CONDITION					
		THIN		NORMAL		OVERWEIGHT	
		grams	pouch	grams	pouch	grams	pouch
ADULT CAT	2	170	2	140	1 +1/2	110	1 +1/2
	2.5	195	2 +1/2	165	2	130	1 +1/2
	3	225	2 +1/2	185	2	150	2
	3.5	250	3	210	2 +1/2	165	2
	4	275	3	230	2 +1/2	185	2
	4.5	300	3 +1/2	250	3	200	2 +1/2
	5	325	4	270	3	215	2 +1/2
	5.5	345	4	290	3 +1/2	230	2 +1/2
	6	365	4 +1/2	305	3 +1/2	245	3
	6.5	390	4 +1/2	325	4	260	3
	7	410	5	340	4	275	3
	7.5	430	5	360	4	285	3 +1/2
	8	450	5 +1/2	375	4 +1/2	300	3 +1/2
	8.5	470	5 +1/2	390	4 +1/2	315	3 +1/2
	9	490	6	410	5	325	4
	9.5	510	6	425	5	340	4
	10	530	6	440	5	350	4

One pouch is equivalent to 18 g of Urinary S/O dry product.

Urinary S/O is a complete dietetic feed for cats, formulated to dissolve struvite stones and reduce their recurrence through its urine acidifying properties.

**RECOMMENDATIONS:** It is recommended that a veterinarian's opinion be sought before use. Feed Urinary S/O for 5 to 12 weeks for the dissolution of struvite stones and up to 6 months for the reduction of struvite stone recurrence. The cat should be re-evaluated regularly by the veterinarian to assess the relevance of a lifelong recommendation. Water should be available at all times.

**NOTE:** In older cats, renal function should be checked before recommending a Urinary S/O diet. The use of Urinary Ageing 7+ Bladder Comfort should be considered in older cats.

## LOWER URINARY TRACT DISEASE

### RECOMMENDED FOR CASES OF

- Struvite uroliths: dissolution and management of recurrence
- Calcium oxalate uroliths: management of recurrence
- Feline idiopathic cystitis (FIC)

### NOT RECOMMENDED IN CASE OF

- Chronic Kidney Disease (CKD)
- Heart disease (when sodium restriction is sought)
- Concurrent use of urine-acidifying medication
- Growth, gestation/lactation



### IDIOPATHIC CYSTITIS

Thanks to the high moisture content of wet food, Urinary S/O Feline supports urine dilution.



### STRUVITE DISSOLUTION

Urinary S/O Feline helps dissolve struvite uroliths.



### LOW RSS

Undersaturated urine creates a urinary environment unfavourable to the development of struvite and calcium oxalate crystals.



### URINE DILUTION

Diluting urine decreases the urinary concentration of struvites and calcium oxalates. The larger urinary volume also supports regular bladder wash-out.



**[S/O® INDEX]** This diet is formulated to promote a healthy urinary environment.

### KEY VALUES

	Per 1000 kcat	Per 100g as fed
Moisture	948 g	79.60 g
Protein	132 g	11.10 g
Fat Content	41.67 g	3.50 g
Crude Fibre	10.71 g	0.90 g
Ash	28.57 g	2.40 g
Carbohydrate [NFE]	29.76 g	2.50 g
Dietary Fibre	17.86 g	1.50 g
Omega-6	9.52 g	0.80 g
Omega-3	2.38 g	0.20 g
EPA + DHA	1.37 g	0.12 g
Calcium	3.69 g	0.31 g
Phosphorus	3.33 g	0.28 g
Sodium	3.81 g	0.32 g
Magnesium	0.21 g	0.02 g
Potassium	2.62 g	0.22 g
Copper	4.88 mg	0.41 mg
Chloride	4.64 g	0.39 g
Vitamin A	7738 IU	650 IU
Vitamin D	107 IU	9 IU
Metabolisable energy*		84 Kcal

\*Calculated according to NRC 2006 CF.

### COMPOSITION

Poultry by-products, pork by-products, corn starch mixture, minerals, powder cellulose, pork blood products, dried tomato pulp, fish oil, marigold meal.

### ADDITIVES (KG)\*\*

Nutritional additives: Vitamin D3: 30 IU, Iron [3b103]: 3 mg, Iodine [3b202]: 0.26 mg, Copper [3b405, 3b406]: 2 mg, Manganese [3b502, 3b503, 3b504]: 1 mg, Zinc [3b603, 3b605, 3b606]: 10 mg - Technological additives: Sodium bisulphate: 0.5 g.

\*\*NOTE: Values reflect only levels added to the formula, not those naturally occurring in components of the diet.

## URINARY S/O MODERATE CALORIE



Available in  
1.5, 3.5,  
7 and 9 kg bags

#### RECOMMENDED DAILY INTAKE & FEEDING DURATION

	CAT'S WEIGHT (kg)	BODY CONDITION	
		NORMAL	OVERWEIGHT
		grams	grams
ADULT CAT	2	36	29
	2.5	42	34
	3	48	39
	3.5	54	43
	4	59	47
	4.5	64	51
	5	69	55
	5.5	74	59
	6	79	63
	6.5	84	67
	7	88	70
	7.5	92	74
	8	97	77
	8.5	101	81
	9	105	84
	9.5	109	87
	10	113	91

Urinary S/O Moderate Calorie is a complete dietetic feed for cats, formulated to dissolve struvite stones and reduce their recurrence through its urine acidifying properties and its low level of magnesium.

**RECOMMENDATIONS:** It is recommended that a veterinarian's opinion be sought before use. Feed Urinary S/O Moderate Calorie for 5 to 12 weeks for the dissolution of struvite stones and up to 6 months for the reduction of struvite stone recurrence. The cat should be re-evaluated regularly by the veterinarian to assess the relevance of a lifelong recommendation. Water should be available at all times.

**NOTE:** In older cats, renal function should be checked before recommending a Urinary S/O diet. The use of Urinary Ageing 7+ Bladder Comfort should be considered in older cats.

**The wet version of Urinary S/O Moderate Calorie is recommended for recurrent idiopathic cystitis.**

## LOWER URINARY TRACT DISEASE

### RECOMMENDED FOR CATS WITH TENDENCY TO BE OVERWEIGHT IN CASES OF

- Struvite uroliths: dissolution and management of recurrence
- Calcium oxalate uroliths: management of recurrence

### NOT RECOMMENDED IN CASE OF

- Chronic Kidney Disease (CKD)
- Heart disease (when sodium restriction is sought)
- Concurrent use of urine-acidifying medication
- Growth, gestation/lactation



### STRUVITE DISSOLUTION

Helps dissolve all types of struvite stones.



### MODERATE CALORIE

Moderate calorie content to help maintain ideal weight.



### URINE DILUTION

Urine dilution makes the urine less liable to form struvite and calcium oxalate stones.



### LOW RSS

Helps lower the concentration of ions contributing to crystal formation.



**[S/O® INDEX]** This diet is formulated to promote a healthy urinary environment.

### KEY VALUES

	Per 1000 kcat	Per 100g as fed
Moisture	15.65 g	5.50 g
Protein	96.76 g	34.00 g
Fat Content	31.30 g	11.00 g
Crude Fibre	19.92 g	7.00 g
Ash	25.33 g	8.90 g
Carbohydrate [NFE]	95.62 g	33.60 g
Dietary Fibre	36.99 g	13.00 g
Omega-6	7.34 g	2.58 g
Omega-3	1.85 g	0.65 g
EPA + DHA	0.94 g	0.33 g
Calcium	2.85 g	1.00 g
Phosphorus	2.56 g	0.90 g
Sodium	3.41 g	1.20 g
Magnesium	0.14 g	0.05 g
Potassium	2.85 g	1.00 g
Copper	4.27 mg	1.50 mg
Chloride	5.69 g	2.00 g
Vitamin A	6545 IU	2300 IU
Vitamin D	228 IU	80 IU
Metabolisable energy*		351.40 Kcal

\*Calculated according to NRC 2006 CF.

### COMPOSITION

Rice, dehydrated poultry proteins, wheat gluten\*\*, maize flour, vegetable fibres, maize gluten, hydrolysed animal proteins, minerals, animal fats, fish oil, soya oil, fructo-oligosaccharides, marigold meal.

\*\* L.I.P.: protein selected for its very high digestibility.

### ADDITIVES (KG)\*\*\*

Nutritional additives: Vitamin A: 21500 IU, Vitamin D3: 800 IU, Iron [3b103]: 38 mg, Iodine [3b201, 3b202]: 3.8 mg, Copper [3b405, 3b406]: 12 mg, Manganese [3b502, 3b504]: 49 mg, Zinc [3b603, 3b605, 3b606]: 141 mg, Selenium [3b801, 3b811, 3b812]: 0.06 mg - Antioxidants.

\*\*\* Values reflect only levels added to the formula, not those naturally occurring in components of the diet.





URINARY S/O

MODERATE CALORIE

# URINARY S/O

## MODERATE CALORIE



MORSELS IN GRAVY

Available in box  
of 12 pouches  
of 85g

### RECOMMENDED DAILY INTAKE & FEEDING DURATION

	CAT'S WEIGHT (kg)	BODY CONDITION			
		NORMAL		OVERWEIGHT	
		grams	pouch	grams	pouch
ADULT CAT	2	160	2	130	1 + 1/2
	2.5	185	2	150	2
	3	215	2 + 1/2	170	2
	3.5	240	3	190	2
	4	260	3	210	2 + 1/2
	4.5	285	3 + 1/2	225	2 + 1/2
	5	305	3 + 1/2	245	3
	5.5	330	4	260	3
	6	350	4	280	3 + 1/2
	6.5	370	4 + 1/2	295	3 + 1/2
	7	390	4 + 1/2	310	3 + 1/2
	7.5	410	5	325	4
	8	430	5	340	4
	8.5	445	5	355	4

One pouch is equivalent to 18 g of Urinary S/O Moderate Calorie dry product.

Urinary S/O Moderate Calorie is a complete dietetic feed for cats, formulated to dissolve struvite stones and reduce their recurrence through its urine acidifying properties and its low level of magnesium. **RECOMMENDATIONS:** It is recommended that a veterinarian's opinion be sought before use. Feed Urinary S/O Moderate Calorie for 5 to 12 weeks for the dissolution of struvite stones and up to 6 months for the reduction of struvite stone recurrence. The cat should be re-evaluated regularly by the veterinarian to assess the relevance of a lifelong recommendation. Water should be available at all times.

**NOTE:** In older cats, renal function should be checked before recommending a Urinary S/O diet. The use of Urinary Ageing 7+ Bladder Comfort should be considered in older cats.

## LOWER URINARY TRACT DISEASE

### RECOMMENDED FOR CASES OF

- Struvite uroliths: dissolution and management of recurrence
- Calcium oxalate uroliths: management of recurrence
- Feline idiopathic cystitis (FIC)

### NOT RECOMMENDED IN CASE OF

- Chronic Kidney Disease (CKD)
- Heart disease (when sodium restriction is sought)
- Concurrent use of urine-acidifying medication
- Growth, gestation/lactation



#### URINE DILUTION

Urine dilution makes the urine less liable to form struvite and calcium oxalate stones.



#### STRUVITE DISSOLUTION

Helps dissolve all types of struvite stones.



#### LOW RSS

Helps lower the concentration of ions contributing to crystal formation.



#### MODERATE CALORIE

Moderate calorie content to help maintain ideal weight.



This diet is formulated to promote a healthy urinary environment.

### KEY VALUES

	Per 1000 kcal	Per 100g as fed
Moisture	1110 g	81.50 g
Protein	129 g	9.50 g
Fat Content	36.78 g	2.70 g
Crude Fibre	14.99 g	1.10 g
Crude Ash	25.89 g	1.90 g
Carbohydrate (NFE)	44.96 g	3.30 g
Dietary Fibre	20.44 g	1.50 g
Omega-6	6.81 g	0.50 g
Omega-3	2.45 g	0.18 g
EPA + DHA	1.16 g	0.09 g
Calcium	3.68 g	0.27 g
Phosphorus	3 g	0.22 g
Sodium	3.68 g	0.27 g
Magnesium	0.14 g	0.01 g
Potassium	2.18 g	0.16 g
Copper	4.63 mg	0.34 mg
Chloride	5.31 g	0.39 g
Vitamin A	8856 IU	650 IU
Vitamin D	341 IU	25 IU
Metabolisable energy*		73.40 Kcal

\*Calculated according to NRC 2006 CF.

### COMPOSITION

Pork by-products, poultry by-products, salmon, pork blood products, wheat flour, powder cellulose, vital wheat gluten, corn starch mixture, minerals, fish oil, marigold meal, yeasts and part thereof.

### ADDITIVES (KG)\*\*

Nutritional additives: Vitamin D3: 70 IU, Iron [3b103]: 2 mg, Iodine [3b202]: 0.27 mg, Copper [3b405, 3b406]: 2.2 mg, Manganese [3b502, 3b503, 3b504]: 0.6 mg, Zinc [3b603, 3b605, 3b606]: 6 mg.

\*\* Values reflect only levels added to the formula, not those naturally occurring in components of the diet.



URINARY S/O

+ BLADDER COMFORT



Available in 2 and 4 kg bags

### RECOMMENDED DAILY INTAKE & FEEDING DURATION

	CAT'S WEIGHT (kg)	BODY CONDITION	
		NORMAL	OVERWEIGHT
		grams	grams
ADULT CAT	2	36	28
	2.5	42	33
	3	48	38
	3.5	53	42
	4	58	47
	4.5	63	51
	5	68	55
	5.5	73	58
	6	78	62
	6.5	82	66
	7	87	69
	7.5	91	73
	8	96	76
	8.5	100	80

ROYAL CANIN® Urinary S/O + Bladder Comfort dry is a complete dietetic feed for cats formulated for the dissolution of struvite stones and the reduction of struvite stone recurrence, and for feline lower urinary tract disease. Urine undersaturating or metastabilising properties for struvite and/or urine acidifying properties.

**RECOMMENDATIONS:** It is recommended that a veterinarian's opinion be sought before use and before extending the period of use. Feed Urinary S/O + Bladder Comfort for 5 to 12 weeks for the dissolution of struvite stones and up to 6 months for the reduction of struvite stone recurrence. Water should be available at all times.

**NOTE:** In older cats, renal function should be checked before recommending a Urinary S/O diet. The use of Urinary S/O + Bladder Comfort is recommended in older cats.

### RECOMMENDED FOR CASES OF

- Management of risk factors for Feline Lower Urinary Tract Disease (FLUTD): tendency to be overweight, insufficient water intake and exposure to increased environmental challenges
- Feline idiopathic cystitis (FIC)
- Struvite uroliths: dissolution and management of recurrence
- Calcium oxalate uroliths: management of recurrence
- Stress related urinary condition

### NOT RECOMMENDED FOR CASES OF

- Chronic Kidney Disease (CKD)
- Heart disease (when sodium restriction is required)
- Concurrent use of urine-acidifying medication
- Growth, gestation/lactation



#### STRUVITE DISSOLUTION

Helps dissolve struvite stones.



#### HYDRATION SUPPORT

This diet encourages cats' natural drinking behaviour, promoting urine dilution thus limiting urine saturation.



#### MODERATE CALORIE

A diet with moderate calorie content supports healthy weight maintenance which can help reduce the risk of feline lower urinary tract disease (FLUTD).



#### BLADDER COMFORT

This diet contains a combination of nutrients to address the leading risk factors of feline lower urinary tract disease including feline idiopathic cystitis (FIC).



This diet is formulated to promote a healthy urinary environment.

### KEY VALUES

	Per 1000 Kcal	Per 100g as fed
Moisture	15.32 g	5.5 g
Protein	94.73 g	34 g
Fat Content	30.65 g	11 g
Crude Fibre	15.32 g	5.5 g
Ash	25.91 g	9.3 g
Carbohydrate (NFE)	96.68 g	34.7 g
Dietary Fibre	36.22 g	13 g
Omega-6	6.46 g	2.32 g
Omega-3	2.95 g	1.06 g
EPA + DHA	1.89 g	0.68 g
Calcium	3.06 g	1.1 g
Phosphorus	2.79 g	1 g
Sodium	3.34 g	1.2 g
Magnesium	0.2 g	0.07 g
Potassium	2.23 g	0.8 g
Copper	4.18 mg	1.5 mg
Chloride	5.29 g	1.9 g
Vitamin A	7941 IU	2850 IU
Vitamin D	195 IU	70 IU
Metabolisable energy*		356.3 Kcal

\*Calculated according to NRC 2006 TDF.

### COMPOSITION

Dehydrated poultry proteins, wheat gluten\*\*, maize, maize flour, wheat, vegetable fibres, hydrolysed animal proteins, rice, maize gluten, minerals, animal fats, chicory pulp, fish oil, psyllium husks and seeds, soya oil, fructo-oligosaccharides, hydrolysed milk protein (0.09%), marigold meal.

\*\*LIP: protein selected for its very high digestibility.

### ADDITIVES (KG)\*\*\*

Nutritional additives: Vitamin A: 2 000 IU, Vitamin D3: 700 IU, Iron [3b103]: 34 mg, Iodine [3b201, 3b202]: 3.4 mg, Copper [3b405, 3b406]: 10 mg, Manganese [3b502, 3b504]: 44 mg, Zinc [3b603, 3b605, 3b606]: 131 mg, Selenium [3b801, 3b811, 3b812]: 0.07 mg - Antioxidants.

\*\*\*Values reflect only levels added to the formula, not those naturally occurring in components of the diet.



URINARY S/O + BLADDER COMFORT



URINARY S/O + BLADDER COMFORT + SATIETY



## URINARY S/O + BLADDER COMFORT + SATIETY



Available in 2 kg bags

### RECOMMENDED DAILY INTAKE & FEEDING DURATION

ADULT CAT	IDEAL WEIGHT (kg)	STAGE			
		WEIGHT LOSS		MAINTENANCE AFTER WEIGHT LOSS	
		grams	cups	grams	Cups
	2	28	2/8	30	3/8
	2.5	33	2/8	35	3/8
	3	38	3/8	40	4/8
	3.5	42	3/8	45	4/8
	4	47	4/8	49	5/8
	4.5	51	4/8	54	5/8
	5	55	4/8	58	6/8
	5.5	58	4/8	62	6/8
	6	62	5/8	66	6/8
	6.5	66	5/8	70	7/8
	7	69	5/8	73	7/8
	7.5	73	6/8	77	7/8
	8	76	6/8	81	1
	8.5	80	6/8	84	1
	9	83	6/8	88	1
	9.5	86	6/8	91	1 + 1/8
	10	89	7/8	95	1 + 1/8

Weight loss: Individual target weight is determined and then feeding amount established. The recommended daily intake is based on the cat's ideal weight. The starting ration may need to be adjusted during follow-up visits, based on each individual's response and the rate of weight loss. An objective of losing 0.5-2% bodyweight per week is generally achievable over the first 3 months, and of 0.5-1% afterwards. Maintenance after weight loss: Energy requirements to maintain stable body weight after weight loss are lower than those of cats that have never had obesity. Based on current scientific knowledge, our maintenance after weight loss rations are calculated by increasing the weight loss ration by about 10%. It is recommended to divide the daily intake into several meals and have fresh water available at all times. ROYAL CANIN® URINARY S/O + BLADDER COMFORT + SATIETY dry is a complete dietetic feed for cats formulated for the dissolution of struvite stones, the reduction of struvite stone recurrence, for feline lower urinary tract disease and for the reduction of excessive body weight. Urine undersaturating or metastabilising properties for struvite and/or urine acidifying properties. RECOMMENDATIONS: your advice as a veterinarian is necessary before use and before extending the period of use of this diet. Feed this diet to 12 weeks for the dissolution of struvite stones and initially up to 6 months before recheck for the reduction of struvite stone recurrence. Feed until the target body weight is achieved and after if required to maintain target body weight. For cats, a transition period when starting the diet is advised. For an efficient weight loss or ideal weight maintenance, the recommended daily energy intake should not be exceeded. The diet may be continued for the cat's lifetime with vet follow-up.

### RECOMMENDED FOR CASES OF

#### Urinary disorders:

- Feline Idiopathic Cystitis (FIC)
- Struvite uroliths and urethral plugs: dissolution and management of recurrence
- Calcium oxalate uroliths: management of recurrence
- Stress or anxiety-related urinary conditions
- Management of risk factors for Feline Lower Urinary Tract Disease (FLUTD) : overweight, insufficient water intake and exposure to increased environmental challenges

#### + Weight management:

- Overweight/obesity: weight loss and maintenance after weight loss
- Stabilised diabetes mellitus with concomitant overweight/obesity
- Hyperlipidaemia with concomitant overweight/obesity

### NOT RECOMMENDED FOR CASES OF

- Growth, gestation or lactation
- Azotaemic Chronic Kidney Disease (IRIS stage 2 to 4)
- Concurrent use of urine-acidifying drugs
- Congestive Heart Failure / Cardiac Disease [when sodium restriction is sought]
- Conditions where high fibre content is not recommended [eg: chronic constipation, severe arthritis making defecation difficult]



#### STRUVITE DISSOLUTION

Helps dissolve struvite stones



#### HYDRATION SUPPORT

This diet encourages cats' natural drinking behaviour. It promotes urine dilution thus limiting urine saturation.



#### BLADDER COMFORT

This diet contains a combination of nutrients to address the leading risk factors of FLUTD including FIC.



#### WEIGHT MANAGEMENT

Formulated to support safe weight loss, control begging and help limit weight regain.



This diet is formulated to promote a healthy urinary environment.

### KEY VALUES

	Per 1000 kcal	Per 100g as fed
Moisture	17.59 g	5.50 g
Protein	109 g	34 g
Fat content	28.79 g	9 g
Crude Fibre	46.71 g	14.60 g
Ash	26.23 g	8.20 g
Carbohydrate (NFE)	90.85 g	28.4 g
Dietary Fibre	78.37 g	24.50 g
Omega-6 fatty acids	5.76 g	1.80 g
Omega-3 fatty acids	1.79 g	0.56 g
EPA + DHA	1.02 g	0.32 g
Calcium	2.85 g	0.89 g
Phosphorus	2.56 g	0.80 g
Potassium	2.59 g	0.81 g
Sodium	3.61 g	1.13 g
Chloride	6.05 g	1.89 g
Magnesium	0.16 g	0.05 g
Vitamin A	8890 IU	2520 IU
Vitamin D	289 IU	100 IU
Metabolisable energy*		312.60 kcal

\*Calculated according to NRC 2006 CF.

Sulphur 0.71%


### COMPOSITION

Dehydrated poultry proteins, vegetable fibres, wheat gluten, manioc (tapioca), wheat flour, hydrolysed animal proteins, minerals, animal fats, chicory pulp, maize gluten, fish oil, maize flour, psyllium husks and seeds, marigold meal, hydrolysed milk protein, glucosamine from fermentation, hydrolysed cartilage (source of chondroitin).

### ADDITIVES (KG)\*\*

Nutritional additives: Vitamin A: 25200 IU, Vitamin D3: 1000 IU, Iron (3b103): 30 mg, Iodine (3b201, 3b202): 3.1 mg, Copper (3b405, 3b406): 9 mg, Manganese (3b502, 3b504): 40 mg, Zinc (3b603, 3b605, 3b606): 95 mg, Selenium (3b607, 3b611, 3b612): 0.04 mg - Antioxidants

\*\* Values reflect only levels added to the formula, not those naturally occurring in components of the diet.



## URINARY S/O + HYPOALLERGENIC



Available in 2 kg bags

### RECOMMENDED DAILY INTAKE & FEEDING DURATION

ADULT CAT	CAT'S WEIGHT (kg)	BODY CONDITION		
		THIN	NORMAL	OVERWEIGHT
		grams	grams	grams
	2	40	34	27
	2.5	47	39	32
	3	54	45	36
	3.5	60	50	40
	4	66	55	44
	4.5	72	60	48
	5	77	65	52
	5.5	83	69	55
	6	88	74	59
	6.5	93	78	62
	7	98	82	66
	7.5	103	86	69
	8	108	90	72
	8.5	113	94	75
	9	118	98	78
	9.5	122	102	81
	10	127	106	84

ROYAL CANIN® Urinary S/O + Hypoallergenic dry is a complete dietetic feed for cats formulated for the dissolution of struvite stones, the reduction of struvite stone recurrence, and for the reduction of ingredient and nutrient intolerances. Urine undersaturating or metastabilising properties for struvite and/or urine acidifying properties. Hydrolysed protein sources and selected carbohydrate sources.

RECOMMENDATIONS: It is recommended that advice from a veterinarian be sought before use and before extending the period of use. Feed URINARY Urinary S/O + Hypoallergenic dry for 5 to 12 weeks for the dissolution of struvite stones, initially up to 6 months for the reduction of struvite stone recurrence, and 3 to 8 weeks for the reduction of ingredient and nutrient intolerances: if signs of intolerance disappear this feed can be used initially up to one year.

### RECOMMENDED FOR CASES OF

#### Urinary disorders:

- Struvite uroliths: dissolution and management of recurrence
- Calcium oxalate uroliths: management of recurrence

+

#### Dermatological/GI conditions:

- Adverse Food Reactions (AFR)
- Feline Atopic Skin Syndrome (FASS)
- Chronic diarrhoea
- Inflammatory Bowel Disease (IBD)
- Exocrine Pancreatic Insufficiency (EPI)

### NOT RECOMMENDED FOR CASES OF

- Chronic Kidney Disease (CKD)
- Concurrent use of urine-acidifying medication
- Growth, gestation/lactation
- Cardiac disease (when sodium restriction is required) / Heart failure



#### URINE DILUTION

Urine dilution makes the urine less liable to form crystals.



#### LOW RSS

A low concentration of ions helps the dissolution and limits the formation of common urinary stones.



#### HYDROLYSED PROTEIN

Hydrolysed protein with low molecular weight to ensure the food is hypoallergenic.



#### SKIN BARRIER

Formulated to support the skin's natural protective barrier for optimal skin health.



This diet is formulated to promote a healthy urinary environment.

### KEY VALUES

	Per 1000 Kcal	Per 100g as fed
Moisture	14.58 g	5.5 g
Protein	71.58 g	27 g
Fat Content	42.42 g	16 g
Crude Fibre	14.32 g	5.4 g
Ash	22.8 g	8.6 g
Carbohydrate (NFE)	99.42 g	37.5 g
Dietary Fibre	26.51 g	10 g
Omega-6	11.72 g	4.42 g
Omega-3	2.52 g	0.95 g
EPA + DHA	1.03 g	0.39 g
Calcium	1.86 g	0.7 g
Phosphorus	1.64 g	0.62 g
Potassium	2.12 g	0.8 g
Sodium	3.45 g	1.3 g
Chloride	5.04 g	1.9 g
Magnesium	0.11 g	0.04 g
Copper	3.98 mg	1.5 mg

Metabolisable energy\*

377.2 Kcal

\*Calculated according to NRC 2006 CF.

Essential fatty acids (Linoleic acid: 4.26%, Arachidonic acid: 0.05%).

Sulphur 0.9%.

### COMPOSITION

Rice flour, hydrolysed soya protein isolate, animal fats, vegetable fibres, rice, minerals, hydrolysed poultry liver, chicory pulp, soya oil, fish oil, fructo-oligosaccharides, borage oil, marigold meal.

Protein sources: hydrolysed soya protein isolate (24.8%), hydrolysed poultry liver (3.5%). Carbohydrate sources: rice flour (33.5%), rice (6.4%).

### ADDITIVES (KG)\*\*

Nutritional additives: Vitamin A: 30000 IU, Vitamin D3: 800 IU, Iron (3b103): 39 mg, Iodine (3b201, 3b202): 3.5 mg, Copper (3b405, 3b406): 15 mg, Manganese (3b502, 3b504): 52 mg, Zinc (3b603, 3b605, 3b606): 153 mg, Selenium (3b607, 3b611, 3b612): 0.25 mg - Technological additives: Clnoptilolite of sedimentary origin: 5 g, Sodium bisulphate: 5 g - Antioxidants.

\*\* Values reflect only levels added to the formula, not those naturally occurring in components of the diet.



URINARY S/O + HYPOALLERGENIC

URINARY S/O  
+  
BLADDER COMFORTAvailable in **box of 12 pouches of 85 g**

## RECOMMENDED DAILY INTAKE &amp; FEEDING DURATION

	CAT'S WEIGHT (kg)	BODY CONDITION			
		NORMAL		OVERWEIGHT	
		grams	pouch	grams	pouch
ADULT CAT	2	160	2	125	1 + 1/2
	2.5	185	2	150	2
	3	210	2 + 1/2	170	2
	3.5	235	3	185	2
	4	260	3	205	2 + 1/2
	4.5	280	3 + 1/2	225	2 + 1/2
	5	300	3 + 1/2	240	3
	5.5	325	4	260	3
	6	345	4	275	3
	6.5	365	4 + 1/2	290	3 + 1/2
	7	385	4 + 1/2	305	3 + 1/2
	7.5	405	5	320	4
	8	420	5	335	4
	8.5	440	5	350	4
	9	460	5 + 1/2	365	4 + 1/2
	9.5	475	5 + 1/2	380	4 + 1/2
	10	495	6	395	4 + 1/2

One pouch is equivalent to 18 g of Urinary S/O + Bladder Comfort dry product.

ROYAL CANIN® Urinary S/O + Bladder Comfort morsels in gravy is a complete dietetic feed for cats formulated for the dissolution of struvite stones and the reduction of struvite stone recurrence, and for feline lower urinary tract disease. Urine undersaturating or metastabilising properties for struvite and/or urine acidifying properties.

RECOMMENDATIONS: It is recommended that a veterinarian's opinion be sought before use and before extending the period of use. Feed Urinary S/O + Bladder Comfort for 5 to 12 weeks for the dissolution of struvite stones and up to 6 months for the reduction of struvite stone recurrence. Water should be available at all times.

**NOTE: In older cats, renal function should be checked before recommending a Urinary S/O diet. The use of Urinary S/O Ageing 7+ + Bladder Comfort is recommended in older cats.**

## RECOMMENDED FOR CASES OF

- Management of risk factors for Feline Lower Urinary Tract Disease (FLUTD): tendency to be overweight, insufficient water intake and exposure to increased environmental challenges
- Feline idiopathic cystitis (FIC)
- Struvite uroliths: dissolution and management of recurrence
- Calcium oxalate uroliths: management of recurrence
- Stress related urinary conditions

## NOT RECOMMENDED FOR CASES OF

- Chronic Kidney Disease (CKD)
- Heart disease [when sodium restriction is required]
- Concurrent use of urine-acidifying medication
- Growth, gestation/lactation



## STRUVITE DISSOLUTION

Helps dissolve struvite stones.



## HYDRATION SUPPORT

This diet encourages cats' natural drinking behaviour, promoting urine dilution thus limiting urine saturation.



## MODERATE CALORIE

A diet with moderate calorie content supports healthy weight maintenance which can help reduce the risk of feline lower urinary tract disease (FLUTD).



## BLADDER COMFORT

This diet contains a combination of nutrients to address the leading risk factors of feline lower urinary tract disease including feline idiopathic cystitis (FIC).



This diet is formulated to promote a healthy urinary environment.

## KEY VALUES

	Per 1000 Kcal	Per 100g as fed
Moisture	1081 g	81.5 g
Protein	106 g	8 g
Fat Content	43.77 g	3.3 g
Crude Fibre	15.92 g	1.2 g
Ash	22.55 g	1.7 g
Carbohydrate [NFE]	57.03 g	4.3 g
Dietary Fibre	22.5 g	1.7 g
Omega-6	9.28 g	0.7 g
Omega-3	3.58 g	0.27 g
EPA + DHA	2.12 g	0.16 g
Calcium	1.99 g	0.15 g
Phosphorus	1.72 g	0.13 g
Sodium	3.71 g	0.28 g
Magnesium	0.13 g	0.01 g
Potassium	1.99 g	0.15 g
Copper	3.58 mg	0.27 mg
Chloride	5.31 g	0.4 g
Vitamin A	19629 IU	1480 IU
Vitamin D	292 IU	22 IU
Metabolisable energy*		75.40 Kcal

\*Calculated according to NRC 2006 CF.

## COMPOSITION

Poultry by-products, pork by-products, salmon, maize flour, pork blood products, powder cellulose, maize starch, minerals, vital wheat gluten, fish oil, hydrolysed milk protein (0.03%), marigold meal, yeasts and part thereof.

## ADDITIVES (KG)\*\*

Nutritional additives: Vitamin D3: 50 IU, Taurine: 1 g, Iron [3b103]: 4 mg, Iodine [3b202]: 0.21 mg, Copper [3b405, 3b406]: 1.4 mg, Manganese [3b502, 3b503, 3b504]: 1.2 mg, Zinc [3b603, 3b605, 3b606]: 12 mg - Technological additives: Clinoptilolite of sedimentary origin: 0.4 g.

\*\*Values reflect only levels added to the formula, not those naturally occurring in components of the diet.

URINARY S/O  
AGEING<sup>7+</sup>  
+  
BLADDER COMFORTAvailable in **2 kg bags**

## RECOMMENDED DAILY INTAKE &amp; FEEDING DURATION

	CAT'S WEIGHT (kg)	BODY CONDITION		
		THIN	NORMAL	OVERWEIGHT
		grams	grams	grams
ADULT CAT	2	41	34	27
	2.5	48	40	32
	3	55	46	37
	3.5	61	51	41
	4	67	56	45
	4.5	73	61	49
	5	79	66	53
	5.5	84	70	56
	6	90	75	60
	6.5	95	79	63
	7	100	84	67
	7.5	105	88	70
	8	110	92	73
	8.5	115	96	77
	9	120	100	80
	9.5	124	104	83
	10	129	108	86

ROYAL CANIN® Urinary S/O Ageing 7+ + Bladder Comfort dry is a complete dietetic feed for cats formulated for the dissolution of struvite stones and the reduction of struvite stone recurrence, and for feline lower urinary tract disease. Urine undersaturating or metastabilising properties for struvite and/or urine acidifying properties.

RECOMMENDATIONS: It is recommended that a veterinarian's opinion be sought before use and before extending the period of use. Feed Urinary S/O Ageing 7+ + Bladder Comfort for 5 to 12 weeks for the dissolution of struvite stones and up to 6 months for the reduction of struvite stone recurrence. Water should be available at all times.

RECOMMENDED FOR AGEING CATS  
(**> 7 YEARS OLD**) IN CASE OF

- Management of risk factors for Feline Lower Urinary Tract Disease (FLUTD): tendency to be overweight, insufficient water intake and exposure to increased environmental challenges
- Feline idiopathic cystitis (FIC)
- Struvite uroliths: dissolution and management of recurrence
- Calcium oxalate uroliths: management of recurrence
- Stress related urinary conditions

## NOT RECOMMENDED FOR CASES OF

- Chronic Kidney Disease (CKD)
- Late stage heart disease and/or congestive heart failure
- Concurrent use of urine-acidifying medication
- Growth, gestation/lactation



## STRUVITE DISSOLUTION

Helps dissolve struvite stones.



## HYDRATION SUPPORT

This diet encourages cats' natural drinking behaviour, promoting urine dilution thus limiting urine saturation.



## HEALTHY AGEING

Formulated with an exclusive blend of nutrients to support cats throughout the ageing process and help maintain renal and heart health.



## BLADDER COMFORT

This diet contains a combination of nutrients to address the leading risk factors of feline lower urinary tract disease including feline idiopathic cystitis (FIC).



This diet is formulated to promote a healthy urinary environment.

## KEY VALUES

	Per 1000 Kcal	Per 100g as fed
Moisture	14.84 g	5.5 g
Protein	86.37 g	32 g
Fat Content	29.69 g	11 g
Crude Fibre	9.72 g	3.6 g
Ash	19.43 g	7.2 g
Carbohydrate [NFE]	110 g	40.7 g
Dietary Fibre	29.69 g	11 g
Omega-6	7.1 g	2.63 g
Omega-3	2.02 g	0.75 g
EPA + DHA	1.21 g	0.45 g
Calcium	1.62 g	0.6 g
Phosphorus	1.62 g	0.6 g
Sodium	1.08 g	0.4 g
Magnesium	0.19 g	0.07 g
Potassium	4.59 g	1.7 g
Copper	4.05 mg	1.5 mg
Chloride	4.59 g	1.7 g
Vitamin A	5938 IU	2200 IU
Vitamin D	189 IU	70 IU
Metabolisable energy*		370.5 Kcal

\*Calculated according to NRC 2006 CF.

## COMPOSITION

Maize, wheat gluten\*\*, wheat, dehydrated poultry proteins, wheat flour, maize gluten, hydrolysed animal proteins, animal fats, minerals, rice, chicory pulp, yeasts products, vegetable fibres, egg powder, fish oil, dried tomato pulp, psyllium husks and seeds, soya oil, fructo-oligosaccharides, algal oil Schizochytrium sp, hydrolysed milk protein (0.09%), glucosamine from fermentation, marigold meal, hydrolysed cartilage.

\*\*LIP: protein selected for its very high digestibility.

## ADDITIVES (KG)\*\*\*


Nutritional additives: Vitamin A: 19500 IU, Vitamin D3: 700 IU, Iron [3b103]: 33 mg, Iodine [3b201, 3b202]: 3.3 mg, Copper [3b405, 3b406]: 10 mg, Manganese [3b502, 3b504]: 42 mg, Zinc [3b603, 3b605, 3b606]: 135 mg, Selenium [3b801, 3b811, 3b812]: 0.07 mg - Antioxidants.

\*\*\*Values reflect only levels added to the formula, not those naturally occurring in components of the diet.





# CANINE NUTRITIONAL MANAGEMENT DECISION TREE



## URINARY S/O



Available in  
2, 7.5 and  
13kg bags

### RECOMMENDED DAILY INTAKE & FEEDING DURATION

ADULT DOG	DOG'S WEIGHT (kg)	BODY CONDITION		
		THIN	NORMAL	OVERWEIGHT
		grams	grams	grams
	2	54	48	41
	4	91	80	69
	6	123	109	94
	8	153	135	116
	10	181	159	138
	15	245	216	187
	20	305	268	232
	25	360	317	274
	30	413	363	314
	35	463	408	352
	40	512	451	389
	45	560	492	425
	50	606	533	460
	55	650	572	494
	60	694	611	528
	70	779	686	592
	80	862	758	655

Urinary S/O is a complete dietetic feed for dogs, formulated to dissolve struvite stones and reduce their recurrence through its urine acidifying properties, its low level of magnesium and its restricted level of protein but of high quality.

**RECOMMENDATIONS:** It is recommended that a veterinarian's opinion be sought before use. Feed Urinary S/O for 5 to 12 weeks for the dissolution of struvite stones and up to 6 months for the reduction of struvite stone recurrence. The dog should be re-evaluated regularly by the veterinarian to assess the relevance of a lifelong recommendation. Water should be available at all times.

**NOTE:** In older dogs, the use of Urinary S/O Ageing 7+ is recommended. It is recommended that renal function be checked before recommending a Urinary S/O diet.

It is recommended to divide the daily intake into 2 meals.

## LOWER URINARY TRACT DISEASE

### RECOMMENDED FOR CASES OF

- Struvite uroliths: dissolution and management of recurrence
- Calcium oxalate uroliths: management of recurrence

### NOT RECOMMENDED IN CASE OF

- Chronic Kidney Disease (CKD)
- Heart disease (when sodium restriction is sought)
- Concurrent use of urine-acidifying medication
- Growth, gestation/lactation



### STRUVITE DISSOLUTION

Helps dissolve all types of struvite stones.



### URINE DILUTION

Urine dilution makes the urine less liable to form struvite and calcium oxalate stones.



### LOW RSS

Helps lower the concentration of ions contributing to crystal formation.



### LOW MAGNESIUM

Reduced level of magnesium, a natural component of struvite crystals.



This diet is formulated to promote a healthy urinary environment.

### KEY VALUES

	Per 1000 Kcal	Per 100g as fed
Moisture	24.48 g	9.50 g
Protein	46.38 g	18.00 g
Fat Content	43.80 g	17.00 g
Crude Fibre	5.67 g	2.20 g
Ash	17.01 g	6.60 g
Carbohydrate (NFE)	120.00 g	46.70 g
Dietary Fibre	14.43 g	5.60 g
Omega-6	9.12 g	3.54 g
Omega-3	1.57 g	0.61 g
EPA + DHA	0.57 g	0.22 g
Calcium	1.29 g	0.50 g
Phosphorus	1.29 g	0.50 g
Sodium	3.09 g	1.20 g
Magnesium	0.13 g	0.05 g
Potassium	2.06 g	0.80 g
Copper	4.38 mg	1.70 mg
Chloride	5.93 g	2.30 g
Vitamin A	4509 IU	1750 IU
Vitamin D	258 IU	100 IU
Metabolisable energy*		388.10 Kcal

\*Calculated according to NRC 2006 CF.

### COMPOSITION

Maize flour, rice, animal fats, dehydrated poultry proteins, maize gluten, minerals, hydrolysed animal proteins, vegetable fibres, soya oil, fish oil, fructo-oligosaccharides, mono- and diglycerides of palmitic and stearic acids esterified with citric acid, marigold meal.

### ADDITIVES (KG)\*\*

Nutritional additives: Vitamin A: 16000 IU, Vitamin D3: 1000 IU, Iron [3b103]: 44 mg, Iodine [3b201, 3b202]: 4.4 mg, Copper [3b405, 3b406]: 14 mg, Manganese [3b502, 3b504]: 57 mg, Zinc [3b603, 3b605, 3b606]: 147 mg, Selenium [3b801, 3b811, 3b812]: 0.06 mg Preservatives - Antioxidants.

\*\* Values reflect only levels added to the formula, not those naturally occurring in components of the diet.



URINARY S/O





## URINARY S/O



LOAF

Available in  
410g can

## RECOMMENDED DAILY INTAKE &amp; FEEDING DURATION

	DOG'S WEIGHT (kg)	BODY CONDITION					
		THIN		NORMAL		OVERWEIGHT	
		grams	can	grams	can	grams	can
ADULT DOG	2	208	1/2	183	1/2	158	1/2
	4	350	3/4	308	3/4	266	3/4
	6	475	1 + 1/4	418	1	361	1
	8	589	1 + 1/2	519	1 + 1/4	448	1
	10	697	1 + 3/4	613	1 + 1/2	529	1 + 1/4
	15	944	2 + 1/4	831	2	718	1 + 3/4
	20	1172	2 + 3/4	1031	2 + 1/2	890	2 + 1/4
	25	1385	3 + 1/2	1219	3	1053	2 + 1/2
	30	1588	3 + 3/4	1397	3 + 1/2	1207	3
	35	1783	4 + 1/4	1569	3 + 3/4	1355	3 + 1/4
	40	1970	4 + 3/4	1734	4 + 1/4	1498	3 + 3/4
	45	2152	5 + 1/4	1894	4 + 1/2	1636	4
	50	2329	5 + 3/4	2050	5	1770	4 + 1/4
	55	2502	6	2202	5 + 1/4	1902	4 + 3/4
	60	2671	6 + 1/2	2350	5 + 3/4	2030	5
	70	2998	7 + 1/4	2638	6 + 1/2	2279	5 + 1/2
	80	3314	8	2916	7	2519	6 + 1/4

One can is equivalent to 99 g of Urinary S/O dry product.

Urinary S/O is a complete dietetic feed for dogs, formulated to dissolve struvite stones and reduce their recurrence through its urine acidifying properties, its low level of magnesium and its restricted level of protein but of high quality.

**RECOMMENDATIONS:** It is recommended that a veterinarian's opinion be sought before use. Feed Urinary S/O for 5 to 12 weeks for the dissolution of struvite stones and up to 6 months for the reduction of struvite stone recurrence. The dog should be re-evaluated regularly by the veterinarian to assess the relevance of a lifelong recommendation. Water should be available at all times.

**NOTE:** In older dogs, the use of Urinary S/O Ageing 7+ is recommended. It is recommended that renal function be checked before recommending a Urinary S/O diet.

It is recommended to divide the daily intake into 2 meals.

## LOWER URINARY TRACT DISEASE

## RECOMMENDED FOR CASES OF

- Struvite uroliths: dissolution and management of recurrence
- Calcium oxalate uroliths: management of recurrence
- Bacterial cystitis: management of secondary struvite crystalluria (in association with adequate antibiotic therapy)

## NOT RECOMMENDED IN CASE OF

- Chronic Kidney Disease (CKD)
- Heart disease (when sodium restriction is sought)
- Concurrent use of urine-acidifying medication
- Growth, gestation/lactation



## STRUVITE DISSOLUTION

Helps dissolve all types of struvite stones.



## URINE DILUTION

Urine dilution makes the urine less liable to form struvite and calcium oxalate stones.



## LOW RSS

Helps lower the concentration of ions contributing to crystal formation.



This diet is formulated to promote a healthy urinary environment.

## KEY VALUES

	Per 1000 Kcal		Per 100g as fed	
	grams	pouch	grams	pouch
Ash	22.04 g		2.10 g	
Moisture	18.00 g		78.0 g	
Protein	83.95 g		8 g	
Fat Content	62.96 g		6 g	
Carbohydrate (NFE)	47.22 g		4.5 g	
Crude Fibre	14.69 g		1.4 g	
Dietary Fibre	25.18 g		2.4 g	
Calcium	3.04 g		0.29 g	
Phosphorus	2.31 g		0.2 g	
Potassium	2.10 g		0.2 g	
Sodium	3.15 g		0.29 g	
Chloride	4.30 g		0.41 g	
Magnesium	0.21 g		0.02 g	
Copper	3.15 mg		0.3 mg	
Zinc	30.43 mg		2.90 mg	
Taurine	0.42 g		0.04 g	
Vitamin A	7975 IU		760 IU	
Vitamin E	198.74 mg		19 mg	
Omega-6	18.89 g		1.8 g	
Omega-3	1.26 g		0.12 g	
EPA + DHA	0.62 g		0.06 g	
Metabolisable energy*			95.3 Kcal	

Calculated according to NRC 2006 CF.

## COMPOSITION

Pork by-products, poultry by-products, maize flour, powder cellulose, corn starch mixture, sunflower oil refined, minerals, pork blood products, fish oil, marigold meal.

## ADDITIVES (KG)\*\*

Nutritional additives: Vitamin D3: 190 IU, Iron (3b103): 5 mg, Iodine (3b202): 0.22 mg, Copper (3b405, 3b406): 1.8 mg, Manganese (3b502, 3b503, 3b504): 1.6 mg, Zinc (3b603, 3b605, 3b606): 16 mg - Technological additives: Sodium bisulphate: 0.55g.

\*\* Values reflect only levels added to the formula, not those naturally occurring in components of the diet.



## URINARY S/O



THIN SLICES IN GRAVY

Available in box  
of 12 pouches  
of 100g

## RECOMMENDED DAILY INTAKE &amp; FEEDING DURATION

	DOG'S WEIGHT (kg)	BODY CONDITION					
		THIN		NORMAL		OVERWEIGHT	
		grams	pouch	grams	pouch	grams	pouch
ADULT DOG	2	207	2	182	2	157	1 + 1/2
	4	348	3 + 1/2	306	3	264	2 + 1/2
	6	471	4 + 1/2	415	4	358	3 + 1/2
	8	585	6	515	5	444	4 + 1/2
	10	691	7	608	6	525	5 + 1/2
	15	973	9 + 1/2	824	8	712	7
	20	1162	11 + 1/2	1023	10	883	9
	25	1374	13 + 1/2	1209	12	1044	10 + 1/2
	30	1576	16	1386	14	1197	12
	35	1769	17 + 1/2	1556	15 + 1/2	1344	13 + 1/2
	40	1955	19 + 1/2	1720	17	1486	15

One pouch is equivalent to 24 g of Urinary S/O Small Dog dry product. Urinary S/O is a complete dietetic feed for dogs, formulated to dissolve struvite stones and reduce their recurrence through its urine acidifying properties, its low level of magnesium and its restricted level of protein but of high quality.

**RECOMMENDATIONS:** It is recommended that a veterinarian's opinion be sought before use. Feed Urinary S/O for 5 to 12 weeks for the dissolution of struvite stones and up to 6 months for the reduction of struvite stone recurrence. The dog should be re-evaluated regularly by the veterinarian to assess the relevance of a lifelong recommendation. Water should be available at all times.

**NOTE:** In older dogs, the use of Urinary S/O Ageing 7+ is recommended. It is recommended that renal function be checked before recommending a Urinary S/O diet.

It is recommended to divide the daily intake into 2 meals.

## LOWER URINARY TRACT DISEASE

## RECOMMENDED FOR CASES OF

- Struvite uroliths: dissolution and management of recurrence
- Calcium oxalate uroliths: management of recurrence
- Bacterial cystitis: management of secondary struvite crystalluria (in association with adequate antibiotic therapy)

## NOT RECOMMENDED IN CASE OF

- Chronic Kidney Disease (CKD)
- Heart disease (when sodium restriction is sought)
- Concurrent use of urine-acidifying medication
- Growth, gestation/lactation



## STRUVITE DISSOLUTION

Helps dissolve all types of struvite stones.



## URINE DILUTION

Urine dilution makes the urine less liable to form struvite and calcium oxalate stones.



## LOW RSS

Helps lower the concentration of ions contributing to crystal formation.



This diet is formulated to promote a healthy urinary environment.

## KEY VALUES

	Per 1000 Kcal		Per 100g as fed	
	grams	pouch	grams	pouch
Moisture	837 g		79 g	
Protein	85.81 g		8.1 g	
Fat Content	59.32 g		5.60 g	
Crude Fibre	10.59 g		1.00 g	
Ash	18.01 g		1.70 g	
Carbohydrate (NFE)	48.73 g		4.60 g	
Dietary Fibre	14.83 g		1.40 g	
Omega-6	22.25 g		2.10 g	
Omega-3	1.17 g		0.11 g	
EPA + DHA	0.64 g		0.06 g	
Calcium	1.91 g		0.18 g	
Phosphorus	1.69 g		0.16 g	
Sodium	2.75 g		0.26 g	
Magnesium	0.15 g		0.01 g	
Potassium	1.59 g		0.15 g	
Copper	3.77 mg		0.25 mg	
Chloride	3.92 g		0.37 g	
Vitamin A	6356 IU		600 IU	
Vitamin D	212 IU		20 IU	
Metabolisable energy*			94.4 Kcal	

\*Calculated according to NRC 2006 CF.

## COMPOSITION

Pork by-products, poultry by-products, wheat flour, sunflower oil refined, corn starch mixture, vital wheat gluten, powder cellulose, pork blood products, minerals, fish oil, marigold meal, yeasts and part thereof.

## ADDITIVES (KG)\*\*

Nutritional additives: Vitamin D3: 170 IU, Iron (3b103): 3 mg, Iodine (3b202): 0.16 mg, Copper (3b405, 3b406): 1.3 mg, Manganese (3b502, 3b503, 3b504): 1 mg, Zinc (3b603, 3b605, 3b606): 10 mg - Technological additives: Sodium bisulphate: 0.9g.

\*\* Values reflect only levels added to the formula, not those naturally occurring in components of the diet.





URINARY S/O

SMALL DOGS



# URINARY S/O

SMALL DOGS



Available in  
1.5, 4 and  
8kg bags

## RECOMMENDED DAILY INTAKE & FEEDING DURATION

ADULT DOG	DOG'S WEIGHT (kg)	BODY CONDITION		
		THIN	NORMAL	OVERWEIGHT
		grams	grams	grams
	2	55	48	42
	2.5	65	57	49
	3	74	65	56
	3.5	83	73	63
	4	92	81	70
	5	109	96	83
	6	125	110	95
	7	140	123	106
	8	155	136	117
	9	169	149	128
	10	183	161	139

Urinary S/O Small Dog is a complete dietetic feed for dogs, formulated to dissolve struvite stones and reduce their recurrence through its urine acidifying properties, its low level of magnesium and its restricted level of protein but of high quality.

**RECOMMENDATIONS:** It is recommended that a veterinarian's opinion be sought before use. Feed Urinary S/O Small Dog for 5 to 12 weeks for the dissolution of struvite stones and up to 6 months for the reduction of struvite stone recurrence. The dog should be re-evaluated regularly by the veterinarian to assess the relevance of a lifelong recommendation. Water should be available at all times.

**NOTE:** In older dogs, the use of Urinary S/O Ageing 7+ is recommended. It is recommended that renal function be checked before recommending a Urinary S/O diet.

It is recommended to divide the daily intake into 2 meals.

## LOWER URINARY TRACT DISEASE

### RECOMMENDED FOR CASES OF

- Struvite uroliths: dissolution and management of recurrence
- Calcium oxalate uroliths: management of recurrence
- Bacterial cystitis: management of secondary struvite crystalluria (in association with adequate antibiotherapy)

### NOT RECOMMENDED IN CASE OF

- Chronic Kidney Disease (CKD)
- Heart disease (when sodium restriction is sought)
- Concurrent use of urine-acidifying medication
- Growth, gestation/lactation



### STRUVITE DISSOLUTION

Helps dissolve all types of struvite stones.



### LOW RSS

Helps lower the concentration of ions contributing to crystal formation.



### URINE DILUTION

Urine dilution makes the urine less liable to form struvite and calcium oxalate stones.



### TARTAR CONTROL

Nutrients that trap the calcium in saliva so reducing tartar build-up.



This diet is formulated to promote a healthy urinary environment.

### KEY VALUES

	Per 1000 Kcal	Per 100g as fed
Moisture	24.73 g	9.50 g
Protein	52.07 g	20.00 g
Fat Content	44.26 g	17.00 g
Crude Fibre	6.25 g	2.40 g
Ash	19.01 g	7.30 g
Carbohydrate [NFE]	114.00 g	43.80 g
Dietary Fibre	14.84 g	5.70 g
Omega-6	9.40 g	3.61 g
Omega-3	1.61 g	0.62 g
EPA + DHA	0.57 g	0.22 g
Calcium	1.87 g	0.72 g
Phosphorus	1.82 g	0.70 g
Sodium	3.12 g	1.20 g
Magnesium	0.13 g	0.05 g
Potassium	2.08 g	0.80 g
Copper	4.69 mg	1.80 mg
Chloride	5.21 g	2.00 g
Vitamin A	4634 IU	1780 IU
Vitamin D	260 IU	100 IU
Metabolisable energy*		384.10 Kcal

\*Calculated according to NRC 2006 CF.

### COMPOSITION

Rice, maize flour, dehydrated poultry proteins, animal fats, maize gluten, hydrolysed animal proteins, minerals (including pentasodium triphosphate (0.35%)), vegetable fibres, egg powder, wheat gluten\*\*, soya oil, fish oil, fructo-oligosaccharides, marigold meal.

\*\* L.I.P.: protein selected for its very high digestibility.

### ADDITIVES (KG)\*\*\*

Nutritional additives: Vitamin A: 16000 IU, Vitamin D3: 1000 IU, Iron [3b103]: 45 mg, Iodine [3b201, 3b202]: 4.5 mg, Copper [3b405, 3b406]: 14 mg, Manganese [3b502, 3b504]: 59 mg, Zinc [3b603, 3b605, 3b606]: 144 mg, Selenium [3b801, 3b811, 3b812]: 0.06 mg Preservatives - Antioxidants.

\*\*\*NOTE: Values reflect only levels added to the formula, not those naturally occurring in components of the diet.



# URINARY S/O

## MODERATE CALORIE



Available in  
1.5, 6.5 and  
12.5 kg bags

## RECOMMENDED DAILY INTAKE & FEEDING DURATION

ADULT DOG	DOG'S WEIGHT (kg)	BODY CONDITION	
		NORMAL	OVERWEIGHT
		grams	grams
	2	49	41
	4	82	69
	6	111	93
	8	137	116
	10	163	137
	15	220	186
	20	273	230
	25	323	272
	30	370	312
	35	416	350
	40	460	387
	45	502	423
	50	543	458
	55	584	492
	60	623	525
	65	662	557
	70	699	589

Urinary S/O Moderate Calorie is a complete dietetic feed for dogs, formulated to dissolve struvite stones and reduce their recurrence through its urine acidifying properties, its low level of magnesium and its restricted level of protein but of high quality.

**RECOMMENDATIONS:** It is recommended that a veterinarian's opinion be sought before use. Feed Urinary S/O Moderate Calorie for 5 to 12 weeks for the dissolution of struvite stones and up to 6 months for the reduction of struvite stone recurrence. The dog should be re-evaluated regularly by the veterinarian to assess the relevance of a lifelong recommendation. Water should be available at all times.

**NOTE:** In older dogs, the use of Urinary S/O Ageing 7+ is recommended. It is recommended that renal function be checked before recommending a Urinary S/O diet.

It is recommended to divide the daily intake into 2 meals.

## LOWER URINARY TRACT DISEASE

### RECOMMENDED FOR CASES OF

- Struvite uroliths: dissolution and management of recurrence
- Calcium oxalate uroliths: management of recurrence
- Bacterial cystitis: management of secondary struvite crystalluria (in association with adequate antibiotherapy)

### NOT RECOMMENDED IN CASE OF

- Chronic Kidney Disease (CKD)
- Heart disease (when sodium restriction is sought)
- Concurrent use of urine-acidifying medication
- Growth, gestation/lactation



### STRUVITE DISSOLUTION

Helps dissolve all types of struvite stones.



### MODERATE CALORIE

Moderate calorie content to help maintain ideal weight.



### URINE DILUTION

Urine dilution makes the urine less liable to form struvite and calcium oxalate stones.



### LOW RSS

Helps lower the concentration of ions contributing to crystal formation.



This diet is formulated to promote a healthy urinary environment.

### KEY VALUES

	Per 1000 Kcal	Per 100g as fed
Moisture	28.90 g	9.50 g
Protein	60.85 g	20.00 g
Fat Content	33.47 g	11.00 g
Crude Fibre	19.77 g	6.50 g
Ash	21.90 g	7.20 g
Carbohydrate [NFE]	139.00 g	45.80 g
Dietary Fibre	37.12 g	12.20 g
Omega-6	7.18 g	2.36 g
Omega-3	1.40 g	0.46 g
EPA + DHA	0.61 g	0.20 g
Calcium	1.83 g	0.60 g
Phosphorus	1.83 g	0.60 g
Sodium	3.65 g	1.20 g
Magnesium	0.24 g	0.08 g
Potassium	2.43 g	0.80 g
Copper	5.17 mg	1.70 mg
Chloride	6.93 g	2.10 g
Vitamin A	6997 IU	2300 IU
Vitamin D	426 IU	140 IU
Metabolisable energy*		328.70 Kcal

\*Calculated according to NRC 2006 CF.

### COMPOSITION

Maize flour, rice, dehydrated poultry proteins, vegetable fibres, maize gluten, animal fats, minerals, hydrolysed animal proteins, chicory pulp, wheat gluten\*\*, soya oil, fish oil, fructo-oligosaccharides, mono- and diglycerides of palmitic and stearic acids esterified with citric acid, marigold meal.

\*\* L.I.P.: protein selected for its very high digestibility.

### ADDITIVES (KG)\*\*\*

Nutritional additives: Vitamin A: 21500 IU, Vitamin D3: 1400 IU, Iron [3b103]: 45 mg, Iodine [3b201, 3b202]: 4.5 mg, Copper [3b405, 3b406]: 14 mg, Manganese [3b502, 3b504]: 59 mg, Zinc [3b603, 3b605, 3b606]: 146 mg, Selenium [3b801, 3b811, 3b812]: 0.11 mg Preservatives - Antioxidants.

\*\*\*NOTE: Values reflect only levels added to the formula, not those naturally occurring in components of the diet.



URINARY S/O

MODERATE CALORIE



URINARY S/O

MODERATE CALORIE

## URINARY S/O MODERATE CALORIE



THIN SLICES IN GRAVY

Available in box  
of 12 pouches  
of 100g

### RECOMMENDED DAILY INTAKE & FEEDING DURATION

ADULT DOG	DOG'S WEIGHT (kg)	BODY CONDITION			
		NORMAL		OVERWEIGHT	
		grams	pouch	grams	pouch
	2	181	2	152	1 + 1/2
	4	304	3	256	2 + 1/2
	6	412	4	347	3 + 1/2
	8	512	5	431	4 + 1/2
	10	605	6	509	5
	15	820	8	691	7
	20	1018	10	857	8 + 1/2
	25	1203	12	1013	10
	30	1379	14	1161	11 + 1/2
	35	1548	15 + 1/2	1304	13
	40	1711	17	1441	14 + 1/2

One pouch is equivalent to 25 g of Urinary S/O Moderate Calorie dry product.

Urinary S/O Moderate Calorie is a complete dietetic feed for dogs, formulated to dissolve struvite stones and reduce their recurrence through its urine acidifying properties, its low level of magnesium and its restricted level of protein but of high quality.

**RECOMMENDATIONS:** It is recommended that a veterinarian's opinion be sought before use. Feed Urinary S/O Moderate Calorie for 5 to 12 weeks for the dissolution of struvite stones and up to 6 months for the reduction of struvite stone recurrence. The dog should be re-evaluated regularly by the veterinarian to assess the relevance of a lifelong recommendation. Water should be available at all times.

**NOTE:** In older dogs, the use of Urinary S/O Ageing 7+ is recommended. It is recommended that renal function be checked before recommending a Urinary S/O diet. It is recommended to divide the daily intake into 2 meals.

## LOWER URINARY TRACT DISEASE

### RECOMMENDED FOR DOGS WITH TENDENCY TO BE OVERWEIGHT IN CASES OF

- Struvite uroliths: dissolution and management of recurrence
- Calcium oxalate uroliths: management of recurrence
- Bacterial cystitis: management of secondary struvite crystalluria (in association with adequate antibiotherapy)

### NOT RECOMMENDED IN CASE OF

- Chronic Kidney Disease (CKD)
- Heart disease (when sodium restriction is sought)
- Concurrent use of urine-acidifying medication
- Growth, gestation/lactation



### STRUVITE DISSOLUTION

Helps dissolve all types of struvite stones.



### MODERATE CALORIE

Moderate calorie content to help maintain ideal weight.



### URINE DILUTION

Urine dilution makes the urine less liable to form struvite and calcium oxalate stones.



### LOW RSS

Helps lower the concentration of ions contributing to crystal formation.



This diet is formulated to promote a healthy urinary environment.

### KEY VALUES

	Per 1000 Kcal	Per 100g as fed
Moisture	956 g	80.10 g
Protein	96.66 g	8.10 g
Fat Content	47.73 g	4 g
Crude Fibre	10.74 g	0.90 g
Ash	20.29 g	1.70 g
Carbohydrate [NFE]	62.05 g	5.20 g
Dietary Fibre	16.71 g	1.40 g
Omega-6	14.32 g	1.20 g
Omega-3	1.07 g	0.09 g
EPA + DHA	0.58 g	0.05 g
Calcium	2.39 g	0.20 g
Phosphorus	1.91 g	0.16 g
Sodium	2.98 g	0.25 g
Magnesium	0.16 g	0.01 g
Potassium	1.79 g	0.15 g
Copper	2.74 mg	0.23 mg
Chloride	4.30 g	0.36 g
Vitamin A	6563 IU	550 IU
Vitamin D	203 IU	17 IU
Metabolisable energy*		83.80 Kcal

\*Calculated according to NRC 2006 CF.

### COMPOSITION

Poultry by-products, pork by-products, wheat flour, corn starch mixture, vital wheat gluten, pork blood products, powder cellulose, minerals, sunflower oil refined, marigold meal, yeasts and part thereof.

### ADDITIVES (KG)\*\*

Nutritional additives: Vitamin D3: 140 IU, Iron [3b103]: 3 mg, Iodine [3b202]: 0.14 mg, Copper [3b405, 3b406]: 1.1 mg, Manganese [3b502, 3b503, 3b504]: 0.9 mg, Zinc [3b603, 3b605, 3b606]: 9 mg - Technological additives: Sodium bisulphate: 0.9g.

\*\* Values reflect only levels added to the formula, not those naturally occurring in components of the diet.



## URINARY S/O AGEING 7+



Available in  
1.5, 3.5 and  
8 kg bags

### RECOMMENDED DAILY INTAKE & FEEDING DURATION

ADULT DOG	DOG'S WEIGHT (kg)	BODY CONDITION		
		THIN	NORMAL	OVERWEIGHT
		grams	grams	grams
	2	55	48	42
	4	92	81	70
	6	125	110	95
	8	155	136	117
	10	183	161	139
	15	248	218	188
	20	307	270	233
	25	363	320	276
	30	416	366	316
	35	467	411	355
	40	517	455	393
	45	564	497	429
	50	611	538	464
	55	656	577	499
	60	700	616	532
	70	786	692	597
	80	869	765	660

Urinary S/O Ageing 7+ is a complete dietetic feed for dogs, formulated to dissolve struvite stones and reduce their recurrence through its urine acidifying properties, its low level of magnesium and its restricted level of protein but of high quality.

**RECOMMENDATIONS:** It is recommended that a veterinarian's opinion be sought before use. Feed Urinary S/O Ageing 7+ for 5 to 12 weeks for the dissolution of struvite stones and up to 6 months for the reduction of struvite stone recurrence. The dog should be re-evaluated regularly by the veterinarian to assess the relevance of a life long recommendation. Water should be available at all times.

It is recommended to divide the daily intake into 2 meals.

## LOWER URINARY TRACT DISEASE

### RECOMMENDED FOR AGEING DOGS (> 7 YEARS OLD) IN CASE OF

- Struvite uroliths: dissolution and management of recurrence
- Calcium oxalate uroliths: management of recurrence
- Bacterial cystitis: management of secondary struvite crystalluria (in association with adequate antibiotherapy)

### NOT RECOMMENDED IN CASE OF

- hyperkalaemia
- Concurrent use of urine-acidifying medication
- Growth, gestation/lactation



### URINARY TRACT HEALTH

Struvite stone dissolution & specifically formulated for up to 80% of all urinary stones.



### AGEING SUPPORT

Support dogs' vitality and cognition with a blend of nutrients and antioxidants. Also supports kidney health.



This diet is formulated to promote a healthy urinary environment.

### KEY VALUES

	Per 1000 Kcal	Per 100g as fed
Moisture	24.69 g	9.50 g
Protein	51.98 g	20.00 g
Fat Content	44.18 g	17.00 g
Crude Fibre	6.24 g	2.40 g
Ash	18.45g	7.10 g
Carbohydrate [NFE]	114.35 g	44.00 g
Dietary Fibre	14.81 g	5.70 g
Omega-6	8.94 g	3.44 g
Omega-3	2.21 g	0.85 g
EPA + DHA	1.07 g	0.41 g
Calcium	1.82 g	0.70 g
Phosphorus	1.56 g	0.60 g
Sodium	1.04 g	0.40 g
Magnesium	0.23 g	0.09 g
Potassium	4.42 g	1.70 g
Copper	4.68 mg	1.80 mg
Chloride	3.90 g	1.50 g
Vitamin A	4638.77 IU	1785.00 IU
Vitamin D	265.85 IU	102.30 IU
Metabolisable energy*		384.80 Kcal

\*Calculated according to NRC 2006 CF.

### COMPOSITION

Rice, maize flour, dehydrated poultry proteins, animal fats, wheat flour, hydrolysed animal proteins, minerals, maize gluten, cellulose powder, wheat gluten\*\*, egg powder, fish oil, soya oil, dried tomato pulp [source of lycopene], fructo-oligosaccharides, glucosamine from fermentation, marigold meal, hydrolysed cartilage [source of chondroitin].

\*\* L.I.P.: protein selected for its very high digestibility.

### ADDITIVES (KG)\*\*\*

Nutritional additives: Vitamin A: 16000 IU, Vitamin D3: 1000 IU, Iron [3b103]: 63 mg, Iodine [3b201, 3b202]: 5.5 mg, Copper [3b405, 3b406]: 15 mg, Manganese [3b502, 3b504]: 69 mg, Zinc [3b603, 3b605, 3b606]: 147 mg, Selenium [3b801, 3b811, 3b812]: 0.12 mg - Preservatives - Antioxidants.

\*\*\* NOTE: Values reflect only levels added to the formula, not those naturally occurring in components of the diet.



URINARY S/O

AGEING 7+





URINARY S/O

AGEING 7+

# URINARY S/O AGEING 7+



LOAF

Available in box of 12 pouches of 85g

## RECOMMENDED DAILY INTAKE & FEEDING DURATION

ADULT DOG	DOG'S WEIGHT (kg)	BODY CONDITION					
		THIN		NORMAL		OVERWEIGHT	
		grams	pouch	grams	pouch	grams	pouch
	2	203	2 + 1/2	178	2	154	2
	4	341	4	300	3 + 1/2	259	3
	6	462	5 + 1/2	406	5	351	4
	8	573	6 + 1/2	504	6	435	5
	10	677	8	596	7	515	6
	15	918	11	808	9 + 1/2	698	8
	20	1139	13 + 1/2	1002	12	866	10
	25	1346	16	1185	14	1023	12
	30	1544	18	1358	16	1173	14
	35	1733	20 + 1/2	1525	18	1317	15 + 1/2
	40	1915	22 + 1/2	1686	20	1456	17

One pouch is equivalent to 22 g of Urinary S/O Ageing 7+ DOG dry product.

Urinary S/O Ageing 7+ is a complete dietetic feed for dogs, formulated to dissolve struvite stones and reduce their recurrence through its urine acidifying properties, its low level of magnesium and its restricted level of protein but of high quality.

RECOMMENDATIONS: It is recommended that a veterinarian's opinion be sought before use. Feed Urinary S/O Ageing 7+ for 5 to 12 weeks for the dissolution of struvite stones and up to 6 months for the reduction of struvite stone recurrence. The dog should be re-evaluated regularly by the veterinarian to assess the relevance of a life long recommendation.

Water should be available at all times.

It is recommended to divide the daily intake into 2 meals.

## LOWER URINARY TRACT DISEASE

### RECOMMENDED FOR AGEING DOGS (> 7 YEARS OLD) IN CASE OF

- Struvite uroliths: dissolution and management of recurrence
- Calcium oxalate uroliths: management of recurrence
- Bacterial cystitis: management of secondary struvite crystalluria (in association with adequate antibiotherapy)

### NOT RECOMMENDED IN CASE OF

- hyperkalaemia
- Concurrent use of urine-acidifying medication
- Growth, gestation/lactation



### URINARY TRACT HEALTH

Struvite stone dissolution & specifically formulated for up to 80% of all urinary stones.



### AGEING SUPPORT

Support dogs' vitality and cognition with a blend of nutrients & antioxidants. Also supports kidney health.



This diet is formulated to promote a healthy urinary environment.

### KEY VALUES

	Per 1000 Kcal	Per 100g as fed
Moisture	787 g	77.40 g
Protein	88.14 g	8.70 g
Fat Content	59.96 g	5.90 g
Crude Fibre	13.21 g	1.30 g
Ash	19.31 g	1.90 g
Carbohydrate (NFE)	48.78 g	4.80 g
Dietary Fibre	22.36 g	2.20 g
Omega-6	16.26 g	1.60 g
Omega-3	1.83 g	0.18 g
EPA + DHA	1.12 g	0.11 g
Calcium	1.93 g	0.19 g
Phosphorus	1.42 g	0.14 g
Sodium	1.22 g	0.12 g
Magnesium	0.20 g	0.02 g
Potassium	4.27 g	0.42 g
Copper	3.05 mg	0.30 mg
Chloride	4.37 g	0.43 g
Vitamin A	9146 IU	900 IU
Vitamin D	193 IU	19 IU
Metabolisable energy*		98.40 Kcal

\*Calculated according to NRC 2006 CF.

### COMPOSITION

Pork by-products, poultry by-products, corn starch mixture, powder cellulose, minerals, pork blood products, sunflower oil refined, maize flour, fish oil, dried tomato pulp, marigold meal.

### ADDITIVES (KG)\*\*

Nutritional additives: Vitamin D3: 170 IU, Iron [3b103]: 4 mg, Iodine [3b202]: 0.21 mg, Copper [3b405, 3b406]: 1.7 mg, Manganese [3b502, 3b503, 3b504]: 1.1 mg, Zinc [3b603, 3b605, 3b606]: 11 mg.

\*\* Values reflect only levels added to the formula, not those naturally occurring in components of the diet.

# URINARY U/C



Available in 14kg bags

## RECOMMENDED DAILY INTAKE & FEEDING DURATION

ADULT DOG	DOG'S WEIGHT (kg)	BODY CONDITION		
		THIN	NORMAL	OVERWEIGHT
		grams	grams	grams
	2	55	48	42
	4	92	81	70
	6	125	110	95
	8	155	137	118
	10	184	162	140
	15	249	219	189
	20	309	272	235
	25	365	321	278
	30	419	368	318
	35	470	414	357
	40	520	457	395
	45	568	499	431
	50	614	541	467
	55	660	581	501
	60	704	620	535
	70	791	696	601
	80	874	769	664

Urinary U/C is a complete dietetic feed for dogs, formulated to reduce the formation of urate and cystine stones through its low level of protein and of purines, its high quality protein, its moderate level of sulphur amino acids and its urine alkalisng properties.

RECOMMENDATIONS: Feed Urinary U/C for up to 6 months to reduce the formation of urate stones, for up to 1 year to reduce the formation of cystine stones and for life in the case of irreversible disturbance of uric acid metabolism. It is recommended that a veterinarian's opinion be sought before use or before extending the period of use. The dog should be re-evaluated regularly by the veterinarian to assess the relevance of a lifelong recommendation. Water should be available at all times.

## LOWER URINARY TRACT DISEASE

### RECOMMENDED FOR CASES OF

- Metabolic uroliths (urate, xanthine, cystine): management of recurrence
- Hyperuricosuria
- Management of metabolic uroliths risk in dogs with leishmaniasis treated with xanthine oxidase inhibitors (e.g. allopurinol)\*

\*If presented with a secondary kidney disease (protein losing nephropathy with or without azotaemia) then switch to a Renal diet

### NOT RECOMMENDED IN CASE OF

- Growth, gestation/lactation



### URATE CONTROL

A selection of proteins with low purine content helps reduce the formation of urate urinary stones.



### CYSTINE CONTROL

A limited intake of proteins and of some amino acids helps limit the formation of cystine urinary stones.



### SKIN BARRIER

A complex to support the barrier effect of the skin.



### ANTIOXIDANT COMPLEX

A synergistic complex of antioxidants to help neutralise free radicals.



This diet is formulated to promote a healthy urinary environment.

### KEY VALUES

	Per 1000 Kcal	Per 100g as fed
Moisture	24.82 g	9.50 g
Protein	47.03 g	18.00 g
Fat Content	39.20 g	15.00 g
Crude Fibre	5.75 g	2.20 g
Ash	13.59 g	5.20 g
Carbohydrate (NFE)	131.00 g	50.10 g
Dietary Fibre	14.89 g	5.70 g
Omega-6	7.11 g	2.72 g
Omega-3	1.46 g	0.56 g
EPA + DHA	0.71 g	0.27 g
Calcium	2.09 g	0.80 g
Phosphorus	1.57 g	0.60 g
Sodium	0.78 g	0.30 g
Magnesium	0.13 g	0.05 g
Potassium	2.35 g	0.90 g
Copper	4.44 mg	1.70 mg
Chloride	1.57 g	0.60 g
Methionine + Cysteine	2.30 g	0.88 g
Metabolisable energy*		382.70 Kcal

\*Calculated according to NRC 2006.

### COMPOSITION

Rice, maize flour, wheat flour, animal fats, egg powder, wheat gluten\*\*, minerals, maize gluten, vegetable fibres, fish oil, soya oil, fructo-oligosaccharides, borage oil, marigold meal. Protein sources: egg powder (10%), wheat gluten (6.7%), maize gluten (3%).

\*\* L.I.P.: protein selected for its very high digestibility.

### ADDITIVES (KG)\*\*\*

Nutritional additives: Vitamin A: 30900 IU, Vitamin D3: 825 IU, Iron [3b103]: 46 mg, Iodine [3b201, 3b202]: 4.6 mg, Copper [3b405, 3b406]: 14 mg, Manganese [3b502, 3b504]: 59 mg, Zinc [3b603, 3b605, 3b606]: 150 mg, Selenium [3b801, 3b811, 3b812]: 0.09mg - Preservatives - Antioxidants.

\*\*\* Values reflect only levels added to the formula, not those naturally occurring in components of the diet.







URINARY S/O + SATIETY



Available in 3.5, 9 and 15 kg bags

#### RECOMMENDED DAILY INTAKE & FEEDING DURATION

	IDEAL DOG'S WEIGHT (kg)	BODY CONDITION		
		STARTING WEIGHT LOSS	AFTER 4 WEEKS	MAINTENANCE AFTER WEIGHT LOSS
		grams	grams	grams
		grams	grams	grams
ADULT DOG	2	44 g	38 g	42 g
	4	74 g	64 g	70 g
	6	101 g	86 g	95 g
	8	125 g	107 g	118 g
	10	148 g	127 g	139 g
	15	201 g	172 g	189 g
	20	249 g	213 g	235 g
	25	294 g	252 g	277 g
	30	337 g	289 g	318 g
	35	379 g	324 g	357 g
	40	418 g	359 g	394 g
	45	457 g	392 g	431 g
	50	495 g	424 g	466 g
	55	531 g	455 g	501 g
	60	567 g	486 g	535 g
	70	637 g	546 g	600 g
	80	704 g	603 g	663 g

ROYAL CANIN® Urinary S/O + Satiety dry is a complete dietetic feed for dogs, formulated for the dissolution of struvite stones, the reduction of struvite stone recurrence and for the reduction of excessive body weight. Urine undersaturating or metastabilising properties for struvite and/or Urine acidifying properties.

**RECOMMENDATIONS:** It is recommended that advice from a veterinarian be sought before use and before extending the period of use. Feed Urinary S/O + Satiety dry for 5 to 12 weeks for the dissolution of struvite stones, initially up to 6 months for the reduction of struvite stone recurrence, as well as until the target body weight is achieved and after if required to maintain target body weight. For an efficient weight loss or ideal weight maintenance, the recommended daily energy intake should not be exceeded.

**[S/O® INDEX]** This diet is formulated to promote a healthy urinary environment.

#### RECOMMENDED FOR CASES OF

##### Urinary disorders:

- Struvite uroliths: dissolution and management of recurrence
- Calcium oxalate uroliths: management of recurrence
- Bacterial cystitis: management of secondary struvite crystalluria (in association with adequate antibiotic therapy)

+

##### Weight management:

- Overweight/obesity
- Stabilised Diabetes Mellitus with overweight/obesity
- Hyperlipidaemia with overweight/obesity

#### NOT RECOMMENDED FOR CASES OF

- Chronic Kidney Disease (CKD)
- Cardiac disease (when sodium restriction is recommended) / Heart failure
- Concurrent use of urine-acidifying medication
- Conditions where high fibre content is not recommended (eg: chronic constipation, severe arthritis making defaecation difficult)
- Growth, gestation/lactation



##### URINE DILUTION

Urine dilution makes the urine less liable to form crystals.



##### LOW RSS

Helps lower the concentration of ions contributing to crystal formation.



##### HIGH PROTEIN

A high protein content helps maintain muscle mass throughout the weight loss period.



##### SATIETOGENIC EFFECT

Formulated with a special blend of fibres to promote satiety and reduce the spontaneous consumption of food.

#### KEY VALUES

	Per 1000Kcal	Per 100g as fed
Moisture	35.7 g	9.5 g
Protein	113 g	30 g
Fat Content	33.82 g	9 g
Crude Fibre	61.26 g	16.3 g
Ash	27.43 g	7.3 g
Carbohydrate (NFE)	104.85 g	27.9 g
Dietary Fibre	104 g	27.6 g
Omega-6	7.48 g	1.99 g
Omega-3	2.03 g	0.54 g
EPA + DHA	1.05 g	0.28 g
Calcium	3.38 g	0.9 g
Phosphorus	2.63 g	0.7 g
Potassium	3.01 g	0.8 g
Sodium	3.76 g	1 g
Chloride	6.01 g	1.6 g
Magnesium	0.23 g	0.06 g
Copper	6.39 mg	1.7 mg
Metabolisable energy*		266.1 Kcal

\*Calculated according to NRC 2006 TDF. Sulphur 0.7%.

#### COMPOSITION

Dehydrated poultry proteins, cellulose powder, wheat gluten\*\*, manioc (tapioca), vegetable fibres, hydrolysed animal proteins, wheat, chicory pulp, maize, animal fats, minerals, maize gluten, fish oil, psyllium husks and seeds, fructo-oligosaccharides, soya oil, sodium butyrate, marigold meal, glucosamine from fermentation, hydrolysed cartilage (source of chondroitin).

\*\*LIP protein selected because of its very high digestibility.

#### ADDITIVES (KG)\*\*

Nutritional additives: Vitamin A: 19900 IU, Vitamin D3: 1000 IU, Iron [3b103]: 40 mg, Iodine [3b201, 3b202]: 4 mg, Copper [3b405, 3b406]: 12 mg, Manganese [3b502, 3b504]: 52 mg, Zinc [3b603, 3b605, 3b606]: 137 mg, Selenium [3b801, 3b811, 3b812]: 0.07 mg - Preservatives - Antioxidants.

\*\*\*Values reflect only levels added to the formula, not those naturally occurring in components of the diet.



## URINARY S/O + SATIETY

SMALL DOGS



Available in 1 and 3 kg bags

#### RECOMMENDED DAILY INTAKE & FEEDING DURATION

	IDEAL WEIGHT (kg)	STAGE			
		WEIGHT LOSS		MAINTENANCE AFTER WEIGHT LOSS	
		grams	cups	grams	cups
		grams	cups	grams	cups
ADULT DOG	2	38	3/8	42	4/8
	2.5	45	4/8	49	4/8
	3	51	4/8	57	5/8
	3.5	58	5/8	63	5/8
	4	64	5/8	70	6/8
	4.5	70	6/8	77	7/8
	5	75	6/8	83	7/8
	5.5	81	7/8	89	1
	6	86	7/8	95	1
	6.5	92	1	101	1 + 1/8
	7	97	1	107	1 + 1/8
	7.5	102	1 + 1/8	112	1 + 2/8
	8	107	1 + 1/8	118	1 + 2/8
	8.5	112	1 + 2/8	124	1 + 3/8
	9	117	1 + 2/8	129	1 + 3/8
	9.5	122	1 + 2/8	134	1 + 3/8
	10	127	1 + 3/8	140	1 + 4/8

Weight loss: Individual target weight is determined and then feeding amount established. The recommended daily intake is based on the dog's ideal weight. The starting ration may need to be adjusted during follow-up visits, based on each individual's response and the rate of weight loss. An objective of losing 1-3% bodyweight per week is generally achievable over the first 3 months, and of 0.5-2% afterwards. Maintenance after weight loss: Energy requirements to maintain stable body weight after weight loss are lower than those of dogs that have never had obesity. Based on current scientific knowledge, our maintenance after weight loss rations are calculated by increasing the weight loss ration by about 10%.

It is recommended to divide the daily intake into at least 2 meals and have fresh water available at all times.

ROYAL CANIN® URINARY S/O + SATIETY SMALL DOGS dry is a complete dietetic feed for dogs formulated for the dissolution of struvite stones, the reduction of struvite stone recurrence and for the reduction of excessive body weight. Urine undersaturating or metastabilising properties for struvite and/or urine acidifying properties. RECOMMENDATIONS: your advice as a veterinarian is necessary before use and before extending the period of use of this diet. Feed this diet 5 to 12 weeks for the dissolution of struvite stones and initially up to 6 months before check for the reduction of struvite stone recurrence. Feed until target body weight is achieved and after if required to maintain target body weight. For an efficient weight loss or ideal weight maintenance, the recommended daily energy intake should not be exceeded.

The diet may be continued for the dog's lifetime with vet follow-up.



#### RECOMMENDED FOR CASES OF

##### Urinary disorders:

- Struvite uroliths: dissolution and management of recurrence
- Calcium oxalate uroliths: management of recurrence
- Bacterial cystitis: management of secondary struvite crystalluria (in association with adequate antibiotic therapy)

+

##### Weight management:

- Overweight/obesity: weight loss and maintenance after weight loss
- Stabilised diabetes mellitus with concomitant overweight/obesity
- Hyperlipidaemia with concomitant overweight/obesity

#### NOT RECOMMENDED FOR CASES OF

- Growth, gestation or lactation
- Azotaemic Chronic Kidney Disease (IRIS stage 2 to 4)
- Concurrent use of urine-acidifying drugs
- Congestive Heart Failure/Cardiac Disease (when sodium restriction is sought)
- Conditions where high fibre content is not recommended (eg: chronic constipation, severe arthritis making defaecation difficult)



##### URINE DILUTION

Urine dilution makes the urine less liable to form crystals.



##### LOW RSS

Helps lower the concentration of ions contributing to crystal formation.



##### HIGH PROTEIN

A high protein content helps maintain muscle mass throughout the weight loss period.



##### SATIETOGENIC EFFECT

Formulated with a special blend of fibres to promote satiety and reduce the spontaneous consumption of food.



##### SMALL KIBBLE

Carefully calibrated kibble size to fit small jaws.

**[S/O INDEX]** This diet is formulated to create an environment unfavourable to the formation of struvite and calcium oxalate.

#### KEY VALUES

	Per 1000 kcal	Per 100g as fed
Moisture	37.05 g	9.50 g
Protein	117 g	30 g
Fat content	37.05 g	9.50 g
Crude Fibre	63.57 g	16.30 g
Ash	28.47 g	7.30 g
Carbohydrate (NFE)	107 g	27.40 g
Dietary Fibre	108 g	27.60 g
Omega-6 fatty acids	7.68 g	1.97 g
Omega-3 fatty acids	2.11 g	0.54 g
EPA + DHA	1.09 g	0.28 g
Calcium	3.39 g	0.87 g
Phosphorus	2.73 g	0.70 g
Potassium	3.12 g	0.80 g
Sodium	3.63 g	0.93 g
Chloride	6.05 g	1.55 g
Magnesium	0.23 g	0.06 g
Copper	6.63 mg	1.70 mg
Metabolisable energy*		256.40 kcal

\*Calculated according to NRC 2006 CF.

Sulphur 0.64%

#### COMPOSITION

Dehydrated poultry proteins, cellulose powder, wheat gluten\*\*, manioc (tapioca), vegetable fibres, hydrolysed animal proteins, wheat, chicory pulp, maize, animal fats, minerals, maize gluten, maize flour, fish oil, psyllium husks and seeds, soya oil, sodium butyrate, fructooligosaccharides, marigold meal, glucosamine from fermentation, hydrolysed cartilage (source of chondroitin).

\*\*\*LIP - protein selected for its very high digestibility.

#### ADDITIVES (KG)\*\*

Nutritional additives: Vitamin A: 19900 IU, Vitamin D3: 1000 IU, Iron [3b103]: 40 mg, Iodine [3b201, 3b202]: 4 mg, Copper [3b405, 3b406]: 12 mg, Manganese [3b502, 3b504]: 52 mg, Zinc [3b603, 3b605, 3b606]: 137 mg, Selenium [3b801, 3b811, 3b812]: 0.07 mg - Preservatives - Antioxidants.

\*\*\*Values reflect only levels added to the formula, not those naturally occurring in components of the diet.



URINARY S/O + SATIETY

SMALL DOGS





Available in 3.5, 10 and 15 kg bags

RECOMMENDED DAILY INTAKE & FEEDING DURATION

ADULT DOG	DOG'S WEIGHT (kg)	BODY CONDITION		
		THIN	NORMAL	OVERWEIGHT
		grams	grams	grams
	2	57	50	43
	4	95	84	72
	6	129	114	98
	8	160	141	122
	10	189	167	144
	15	257	226	195
	20	318	280	242
	25	376	331	286
	30	432	380	328
	35	485	426	368
	40	536	471	407
	45	585	515	445
	50	633	557	481
	55	680	598	517
	60	726	639	552
	70	815	717	619
	80	901	793	685

ROYAL CANIN® Urinary S/O + Hypoallergenic dry is a complete dietetic feed for dogs formulated for the dissolution of struvite stones, the reduction of struvite stone recurrence, and for the reduction of ingredient and nutrient intolerances. Urine undersaturating or metastabilising properties for struvite and/or urine acidifying properties. Hydrolysed protein sources and selected carbohydrate sources.

RECOMMENDATIONS: It is recommended that advice from a veterinarian be sought before use and before extending the period of use. Feed Urinary S/O + Hypoallergenic for 5-12 weeks for the dissolution of struvite stones, initially up to 6 months for the reduction of struvite stone recurrence, and 3 to 8 weeks for the reduction of ingredient and nutrient intolerances: if signs of intolerance disappear this feed can be used initially up to one year.

**[S/O®]** This diet is formulated to promote a healthy urinary environment.



**RECOMMENDED FOR CASES OF**  
**Urinary disorders:**

- Struvite uroliths: dissolution and management of recurrence
- Calcium oxalate uroliths: management of recurrence

**•**  
**Dermatological/GI conditions:**

- Adverse Food Reactions (AFR)
- Canine Atopic Dermatitis (CAD)
- Chronic diarrhoea
- Inflammatory Bowel Disease (IBD)
- Exocrine Pancreatic Insufficiency (EPI)

**NOT RECOMMENDED FOR CASES OF**

- Chronic Kidney Disease (CKD)
- Concurrent use of urine-acidifying medication
- Growth, gestation/lactation
- Cardiac disease (when sodium restriction is required) / Heart failure



**URINE DILUTION**  
Urine dilution makes the urine less liable to form crystals.



**LOW RSS**  
A low concentration of ions helps the dissolution and limits the formation of common urinary stones.



**HYDROLYSED PROTEIN**  
Hydrolysed protein with low molecular weight to ensure the food is hypoallergenic.



**SKIN BARRIER**  
Formulated to support the skin's natural protective barrier for optimal skin health.

KEY VALUES	Per 1000Kcal	Per 100g as fed
Moisture	25.59 g	9.5 g
Protein	53.88 g	20 g
Fat Content	40.41 g	15 g
Crude Fibre	6.47 g	2.4 g
Ash	22.36 g	8.3 g
Carbohydrate (NFE)	120.69 g	44.8 g
Dietary Fibre	18.59 g	6.9 g
Omega-6	9.56 g	3.55 g
Omega-3	2.16 g	0.8 g
EPA + DHA	0.97g	0.36 g
Calcium	1.51 g	0.56 g
Phosphorus	1.35 g	0.5 g
Potassium	2.16 g	0.8 g
Sodium	3.37 g	1.25 g
Chloride	4.98 g	1.85 g
Magnesium	0.11 g	0.04 g
Copper	4.04 mg	1.5 mg
Metabolisable energy*		371.2 Kcal

\*Calculated according to NRC 2006 CF. Sulphur 0.6%.

**COMPOSITION**  
Rice flour, hydrolysed soya protein isolate, animal fats, rice, minerals, chicory pulp, soya oil, vegetable fibres, fish oil, fructo-oligosaccharides, sodium butyrate, borage oil, marigold meal.

Protein source: hydrolysed soya protein isolate (17.5%). Carbohydrate sources: rice flour (38.9%), rice (10.1%).

**ADDITIVES (KG)\*\***

Nutritional additives: Vitamin A: 30 500 IU, Vitamin D3: 800 IU, Iron (3b103): 41 mg, Iodine (3b201, 3b202): 3.7 mg, Copper (3b405, 3b406): 15 mg, Manganese (3b502, 3b504): 54 mg, Zinc (3b603, 3b605, 3b606): 160 mg, Selenium (3b801, 3b811, 3b812): 0.26 mg - Technological additives: Clinoptilolite of sedimentary origin: 10 g - Preservatives - Antioxidants.

\*\* Values reflect only levels added to the formula, not those naturally occurring in components of the diet.

COMPATIBLE WITH  
**Treats**  
**URINARY S/O DIETS**



Available in  
6 x 230g bags

RECOMMENDED DAILY INTAKE

Please consult your veterinarian if you have any questions.

Dog's weight	<2 kg	2-5 kg	6-10 kg	11-15 kg	16-20 kg	>20 kg
Max pieces per day	2	4	8	11	14	20

Do not feed more than the maximum number of treats per day indicated in the feeding guidelines. If you are giving treats, it is recommended to adjust the main meal portion size accordingly to maintain your pet's healthy body weight.



**URINARY S/O DIETS**

**TREATS COMPATIBLE WITH URINARY DIETS**  
ROYAL CANIN® Treats were developed by experts in canine nutrition to ensure treating does not compromise the health benefits of your dog's diet.

**COMPATIBLE ROYAL CANIN® DOG DIETS**

- URINARY S/O
- URINARY S/O MODERATE CALORIE
- URINARY S/O SMALL DOG
- URINARY S/O AGEING 7+

**OPTIMISED MINERAL CONTENT**

ANALYTICAL CONSTITUENTS	
Moisture	18%
Protein	18%
Fat Content	5.4%
Crude Fibre	2.4%
Crude Ash	6.8%
Metabolisable energy	2995Kcal/kg

MINERALS (Total)	
Sodium	1.1%

CALORIES PER CHEW	
	4Kcal

**COMPOSITION**  
Rice flour, dehydrated poultry proteins, glycerol, barley flour, minerals, animal fats, wheat gluten, hydrolysed animal proteins, cellulose powder.

**ADDITIVES (KG)**  
Preservatives.





UK

[WWW.ROYALCANIN.COM/UK](http://WWW.ROYALCANIN.COM/UK)

**NUTRITIONAL HELPLINE**

0800 717800

ROI

[WWW.ROYALCANIN.COM/IE](http://WWW.ROYALCANIN.COM/IE)

**NUTRITIONAL HELPLINE**

1800 936074