



AN EXPERT
ALLIANCE FOR
**THE MOST SPECIFIC
SENSITIVITIES**



URINARY S/O + HYPOALLERGENIC



Available in 2 kg bags

RECOMMENDED DAILY INTAKE & FEEDING DURATION

ADULT CAT	CAT'S WEIGHT (kg)	BODY CONDITION		
		THIN	NORMAL	OVERWEIGHT
		grams	grams	grams
	2	40	34	27
	2.5	47	39	32
	3	54	45	36
	3.5	60	50	40
	4	66	55	44
	4.5	72	60	48
	5	77	65	52
	5.5	83	69	55
	6	88	74	59
	6.5	93	78	62
	7	98	82	66
	7.5	103	86	69
	8	108	90	72
	8.5	113	94	75
	9	118	98	78
	9.5	122	102	81
	10	127	106	84

ROYAL CANIN® URINARY S/O + HYPOALLERGENIC dry is a complete dietetic feed for cats formulated for the dissolution of struvite stones, the reduction of struvite stone recurrence, and for the reduction of ingredient and nutrient intolerances. Urine undersaturating or metastabilising properties for struvite and/or Urine acidifying properties. Hydrolysed protein sources and selected carbohydrate sources.

RECOMMENDATIONS: It is recommended that advice from a veterinarian be sought before use and before extending the period of use. Feed URINARY S/O + HYPOALLERGENIC dry for 5 to 12 weeks for the dissolution of struvite stones, initially up to 6 months for the reduction of struvite stone recurrence, and 3 to 8 weeks for the reduction of ingredient and nutrient intolerances: if signs of intolerance disappear this feed can be used initially up to one year.



This diet is formulated to promote a healthy urinary environment.



RECOMMENDED FOR CASES OF

Urinary disorders:

- Struvite uroliths: dissolution and management of recurrence
- Calcium oxalate uroliths: management of recurrence

Dermatological/GI conditions:

- Adverse Food Reactions (AFR)
- Feline Atopic Skin Syndrome (FASS)
- Chronic diarrhoea
- Inflammatory Bowel Disease (IBD)
- Exocrine Pancreatic Insufficiency (EPI)

NOT RECOMMENDED FOR CASES OF

- Chronic Kidney Disease (CKD)
- Concurrent use of urine-acidifying medications
- Growth, gestation/lactation
- Cardiac disease (when sodium restriction is required) / Heart failure



URINE DILUTION

Urine dilution makes the urine less liable to form crystals.



LOW RSS

A low concentration of ions helps the dissolution and limits the formation of common urinary stones.



HYDROLYSED PROTEIN

Hydrolysed protein with low molecular weight to ensure the food is hypoallergenic.



SKIN BARRIER

Formulated to support the skin's natural protective barrier for optimal skin health.

KEY VALUES

	Per 1000Kcal	Per 100g as fed
Moisture	14.58 g	5.5 g
Protein	71.58 g	27 g
Fat content	42.42 g	16 g
Crude Fibre	14.32 g	5.4 g
Ash	22.8 g	8.6 g
Carbohydrate (NFE)	99.42 g	37.5 g
Dietary Fibre	26.51 g	10 g
Omega-6	11.72 g	4.42 g
Omega-3	2.52 g	0.95 g
EPA + DHA	1.03 g	0.39 g
Calcium	1.86 g	0.7 g
Phosphorus	1.64 g	0.62 g
Potassium	2.12 g	0.8 g
Sodium	3.45 g	1.3 g
Chloride	5.04 g	1.9 g
Magnesium	0.11 g	0.04 g
Copper	3.98 mg	1.5 mg
Metabolisable energy*		377.2 Kcal

*Calculated according to NRC 2006 CF.

Essential fatty acids (Linoleic acid: 4.26%, Arachidonic acid: 0.05%). Sulphur 0.9%.

COMPOSITION


Rice flour, hydrolysed soya protein isolate, animal fats, vegetable fibres, rice, minerals, hydrolysed poultry liver, chicory pulp, soya oil, fish oil, fructo-oligosaccharides, borage oil, marigold meal.

Protein sources: hydrolysed soya protein isolate (24.8%), hydrolysed poultry liver (3.5%). Carbohydrate sources: rice flour (33.5%), rice (6.4%).

ADDITIVES (KG)**

Nutritional additives: Vitamin A: 30000 IU, Vitamin D3: 800 IU, Iron [3b103]: 39 mg, Iodine [3b201, 3b202]: 3.5 mg, Copper [3b405, 3b406]: 15 mg, Manganese [3b502, 3b504]: 52 mg, Zinc [3b603, 3b605, 3b606]: 153 mg, Selenium [3b801, 3b811, 3b812]: 0.25 mg - Technological additives: Clinoptilolite of sedimentary origin: 5 g, Sodium bisulphate: 5 g - Antioxidants.

** Values reflect only levels added to the formula, not those naturally occurring in components of the diet.



URINARY S/O + BLADDER COMFORT + SATIETY



Available in 2 kg bags

RECOMMENDED DAILY INTAKE & FEEDING DURATION

ADULT CAT	IDEAL WEIGHT (kg)	STAGE			
		WEIGHT LOSS		MAINTENANCE AFTER WEIGHT LOSS	
		grams	cups	grams	Cups
	2	28	2/8	30	3/8
	2.5	33	2/8	35	3/8
	3	38	3/8	40	4/8
	3.5	42	3/8	45	4/8
	4	47	4/8	49	5/8
	4.5	51	4/8	54	5/8
	5	55	4/8	58	6/8
	5.5	58	4/8	62	6/8
	6	62	5/8	66	6/8
	6.5	66	5/8	70	7/8
	7	69	5/8	73	7/8
	7.5	73	6/8	77	7/8
	8	76	6/8	81	1
	8.5	80	6/8	84	1
	9	83	6/8	88	1
	9.5	86	6/8	91	1 + 1/8
	10	89	7/8	95	1 + 1/8

Weight loss: Individual target weight is determined and then feeding amount established. The recommended daily intake is based on the cat's ideal weight. The starting ration may need to be adjusted during follow-up visits, based on each individual's response and the rate of weight loss. An objective of losing 0.5-2% bodyweight per week is generally achievable over the first 3 months, and of 0.5-1% afterwards. Maintenance after weight loss: Energy requirements to maintain stable bodyweight after weight loss are lower than those of cats that have never had obesity. Based on current scientific knowledge, our maintenance after weight loss rations are calculated by increasing the weight loss ration by about 10%.

It is recommended to divide the daily intake into several meals and have fresh water available at all times.

ROYAL CANIN® URINARY S/O + BLADDER COMFORT + SATIETY dry is a complete dietetic feed for cats formulated for the dissolution of struvite stones, the reduction of struvite stone recurrence, for feline lower urinary tract disease and for the reduction of excessive body weight. Urine undersaturating or metastabilising properties for struvite and/or uric acidifying properties.

RECOMMENDATIONS: your advice as a veterinarian is necessary before use and before extending the period of use of this diet. Feed this diet 5 to 12 weeks for the dissolution of struvite stones and initially up to 6 months before recheck for the reduction of struvite stone recurrence. Feed until the target body weight is achieved and after if required to maintain target body weight. For cats, a transition period when starting the diet is advised.

For an efficient weight loss or ideal weight maintenance, the recommended daily energy intake should not be exceeded. The diet may be continued for the cat's lifetime with vet follow-up.

RECOMMENDED FOR CASES OF

Urinary disorders:

- Feline Idiopathic Cystitis (FIC)
- Struvite uroliths and urethral plugs: dissolution and management of recurrence
- Calcium oxalate uroliths: management of recurrence
- Stress or anxiety-related urinary conditions
- Management of risk factors for Feline Lower Urinary Tract Disease (FLUTD) : overweight, insufficient water intake and exposure to increased environmental challenges



Weight management:

- Overweight/obesity: weight loss and maintenance after weight loss
- Stabilised diabetes mellitus with concomitant overweight/obesity
- Hyperlipidaemia with concomitant overweight/obesity

NOT RECOMMENDED FOR CASES OF

- Growth, gestation or lactation
- Azotaemic Chronic Kidney Disease (IRIS stage 2 to 4)
- Concurrent use of urine-acidifying drugs
- Congestive Heart Failure / Cardiac Disease (when sodium restriction is sought)
- Conditions where high fibre content is not recommended (eg: chronic constipation, severe arthritis making defecation difficult)



STRUVITE DISSOLUTION

Helps dissolve struvite stones



HYDRATION SUPPORT

This diet encourages cats' natural drinking behaviour. It promotes urine dilution thus limiting urine saturation.



BLADDER COMFORT

This diet contains a combination of nutrients to address the leading risk factors of FLUTD including FIC.



WEIGHT MANAGEMENT

Formulated to support safe weight loss, control begging and help limit weight regain.



This diet is formulated to promote a healthy urinary environment.

KEY VALUES

	Per 1000 kcal	Per 100g as fed
Moisture	17.59 g	5.50 g
Protein	109 g	34 g
Fat content	28.79 g	9 g
Crude Fibre	46.71 g	14.60 g
Ash	26.23 g	8.20 g
Carbohydrate (NFE)	90.85 g	28.4 g
Dietary Fibre	78.37 g	24.50 g
Omega-6 fatty acids	5.76 g	1.80 g
Omega-3 fatty acids	1.79 g	0.56 g
EPA + DHA	1.02 g	0.32 g
Calcium	2.85 g	0.89 g
Phosphorus	2.56 g	0.80 g
Potassium	2.59 g	0.81 g
Sodium	3.61 g	1.13 g
Chloride	6.05 g	1.89 g
Magnesium	0.16 g	0.05 g
Vitamin A	8890 IU	2520 IU
Vitamin D	289 IU	100 IU
Metabolisable energy*		312.60 kcal

*Calculated according to NRC 2006 CF.

Sulphur 0.71%

COMPOSITION

Dehydrated poultry proteins, vegetable fibres, wheat gluten, manioc (tapioca), wheat flour, hydrolysed animal proteins, minerals, animal fats, chicory pulp, maize gluten, fish oil, maize flour, psyllium husks and seeds, marigold meal, hydrolysed milk protein, glucosamine from fermentation, hydrolysed cartilage (source of chondroitin).

ADDITIVES (KG)***

Nutritional additives: Vitamin A: 25200 IU, Vitamin D3: 1000 IU, Iron (3b103): 30 mg, Iodine (3b201, 3b202): 3.1 mg, Copper (3b405, 3b406): 9 mg, Manganese (3b502, 3b504): 40 mg, Zinc (3b603, 3b605, 3b606): 95 mg, Selenium (3b801, 3b811, 3b812): 0.04 mg - Antioxidants.

** Values reflect only levels added to the formula, not those naturally occurring in components of the diet.





URINARY S/O + BLADDER COMFORT



Available in 2 and 4 kg bags

RECOMMENDED DAILY INTAKE & FEEDING DURATION

ADULT CAT	CAT'S WEIGHT (kg)	BODY CONDITION	
		NORMAL	OVERWEIGHT
		grams	grams
	2	36	28
	2.5	42	33
	3	48	38
	3.5	53	42
	4	58	47
	4.5	63	51
	5	68	55
	5.5	73	58
	6	78	62
	6.5	82	66
	7	87	69
	7.5	91	73
	8	96	76
	8.5	100	80
	9	104	83
	9.5	108	86
	10	112	89

ROYAL CANIN® MULTIFUNCTION URINARY S/O + BLADDER COMFORT dry is a complete dietetic feed for cats formulated for the dissolution of struvite stones and the reduction of struvite stone recurrence, and for feline lower urinary tract disease. Urine undersaturating or metastabilising properties for struvite and/or urine acidifying properties.

RECOMMENDATIONS: It is recommended that a veterinarian's opinion be sought before use and before extending the period of use. Feed URINARY S/O + BLADDER COMFORT for 5 to 12 weeks for the dissolution of struvite stones and up to 6 months for the reduction of struvite stone recurrence. Water should be available at all times.

NOTE: In older cats renal function should be checked before recommending a Urinary S/O diet. The use of Urinary S/O Ageing + + Bladder Comfort is recommended in older cats.

[S/O INDEX] This diet is formulated to promote a healthy urinary environment.



RECOMMENDED FOR CASES OF

- Management of risk factors for Feline Lower Urinary Tract Disease (FLUTD): tendency to be overweight, insufficient water intake and exposure to increased environmental challenges
 - Feline Idiopathic Cystitis (FIC)
 - Struvite uroliths: dissolution and management of recurrence
 - Calcium oxalate uroliths: management of recurrence
 - Stress related urinary conditions
- ### NOT RECOMMENDED FOR CASES OF
- Chronic Kidney Disease (CKD)
 - Heart disease (when sodium restriction is required)
 - Concurrent use of urine-acidifying medications
 - Growth, gestation/lactation



STRUVITE DISSOLUTION

Helps dissolve struvite stones.



HYDRATION SUPPORT

This diet encourages cats' natural drinking behaviour, promoting urine dilution thus limiting urine saturation.



MODERATE CALORIE

A diet with moderate calorie content supports healthy weight maintenance which can help reduce the risk of feline lower urinary tract disease (FLUTD).



BLADDER COMFORT

This diet contains a combination of nutrients to address the leading risk factors of feline lower urinary tract disease including feline idiopathic cystitis (FIC).

KEY VALUES

	Per 1000Kcal	Per 100g as fed
Moisture	15.44 g	5.5 g
Protein	95.43 g	34 g
Fat content	30.87 g	11 g
Crude Fibre	15.44 g	5.5 g
Ash	26.1 g	9.3 g
Carbohydrate (NFE)	97.39 g	34.7 g
Dietary Fibre	36.49 g	13 g
Omega-6	6.51 g	2.32 g
Omega-3	2.98 g	1.06 g
EPA + DHA	1.91 g	0.68 g
Calcium	3.09 g	1.1 g
Phosphorus	2.81 g	1 g
Sodium	3.37 g	1.2 g
Magnesium	0.2 g	0.07 g
Potassium	2.25 g	0.8 g
Copper	4.21 mg	1.5 mg
Chloride	5.33 g	1.9 g
Vitamin A	7999 IU	2850 IU
Vitamin D	196 IU	70 IU
Metabolisable energy*		356.3 Kcal

*Calculated according to NRC 2006 CF.

COMPOSITION

Dehydrated poultry proteins, wheat gluten**, maize, maize flour, wheat, vegetable fibres, hydrolysed animal proteins, rice, maize gluten, minerals, animal fats, chicory pulp, fish oil, psyllium husks and seeds, soya oil, fructo-oligosaccharides, hydrolysed milk protein (0.09%), marigold meal.

**LIP: protein selected for its very high digestibility.

ADDITIONS (KG)**

Nutritional additives: Vitamin A: 2 000 IU, Vitamin D3: 700 IU, Iron [3b103]: 34 mg, Iodine [3b201, 3b202]: 3.4 mg, Copper [3b405, 3b406]: 10 mg, Manganese [3b502, 3b504]: 44 mg, Zinc [3b603, 3b605, 3b606]: 131 mg, Selenium [3b801, 3b811, 3b812]: 0.07 mg - Antioxidants.

***Values reflect only levels added to the formula, not those naturally occurring in components of the diet.





URINARY S/O AGEING⁷⁺ + BLADDER COMFORT



Available in 2 kg bags

RECOMMENDED DAILY INTAKE & FEEDING DURATION

ADULT CAT	CAT'S WEIGHT (kg)	BODY CONDITION		
		THIN	NORMAL	OVERWEIGHT
		grams	grams	grams
	2	41	34	27
	2.5	48	40	32
	3	55	46	37
	3.5	61	51	41
	4	67	56	45
	4.5	73	61	49
	5	79	66	53
	5.5	84	70	56
	6	90	75	60
	6.5	95	79	63
	7	100	84	67
	7.5	105	88	70
	8	110	92	73
	8.5	115	96	77
	9	120	100	80
	9.5	124	104	83
	10	129	108	86

ROYAL CANIN® MULTIFUNCTION URINARY S/O AGEING⁷⁺ + BLADDER COMFORT dry is a complete dietetic feed for cats formulated for the dissolution of struvite stones and the reduction of struvite stone recurrence, and for feline lower urinary tract disease. Urine undersaturating or metastabilising properties for struvite and/or urine acidifying properties.

RECOMMENDATIONS: it is recommended that a veterinarian's opinion be sought before use and before extending the period of use. Feed URINARY S/O AGEING⁷⁺ + BLADDER COMFORT for 5 to 12 weeks for the dissolution of struvite stones and up to 6 months for the reduction of struvite stone recurrence. Water should be available at all times.



This diet is formulated to promote a healthy urinary environment.



RECOMMENDED FOR AGEING CATS (> 7 YEARS OLD) IN CASE OF

- Management of risk factors for Feline Lower Urinary Tract Disease (FLUTD): tendency to be overweight, insufficient water intake and exposure to increased environmental challenges
- Feline Idiopathic Cystitis (FIC)
- Struvite uroliths: dissolution and management of recurrence
- Calcium oxalate uroliths: management of recurrence
- Stress related urinary conditions
- NOT RECOMMENDED FOR CASES OF**
 - Chronic Kidney Disease (CKD)
 - Late stage heart disease and/or congestive heart failure
 - Concurrent use of urine-acidifying medications
 - Growth, gestation/lactation



STRUVITE DISSOLUTION

Helps dissolve struvite stones.



HYDRATION SUPPORT

This diet encourages cats' natural drinking behaviour, promoting urine dilution thus limiting urine saturation.



HEALTHY AGEING

Formulated with an exclusive blend of nutrients to support cats throughout the ageing process and help maintain renal and heart health.



BLADDER COMFORT

This diet contains a combination of nutrients to address the leading risk factors of feline lower urinary tract disease including feline idiopathic cystitis (FIC).

KEY VALUES

	Per 1000Kcal	Per 100g as fed
Moisture	14.84 g	5.5 g
Protein	86.37 g	32 g
Fat content	29.69 g	11 g
Crude fibre	9.72 g	3.6 g
Ash	19.43 g	7.2 g
Carbohydrate (NFE)	110 g	40.7 g
Dietary fibre	29.69g	11 g
Omega-6	7.1 g	2.63 g
Omega-3	2.02 g	0.75 g
EPA + DHA	1.21 g	0.45 g
Calcium	1.62 g	0.6 g
Phosphorus	1.62 g	0.6 g
Sodium	1.08 g	0.4 g
Magnesium	0.19 g	0.07 g
Potassium	4.59 g	1.7 g
Copper	4.05 mg	1.5 mg
Chloride	4.59 g	1.7 g
Vitamin A	5938 IU	2200 IU
Vitamin D	189 IU	70 IU
Metabolisable energy*		370.5 Kcal

*Calculated according to NRC 2006 CF.

COMPOSITION

Maize, wheat gluten**, wheat, dehydrated poultry proteins, wheat flour, maize gluten, hydrolysed animal proteins, animal fats, minerals, rice, chicory pulp, yeasts products, vegetable fibres, egg powder, fish oil, dried tomato pulp, psyllium husks and seeds, soya oil, fructo-oligosaccharides, algal oil Schizochytrium sp, hydrolysed milk protein [0.09%], glucosamine from fermentation, marigold meal, hydrolysed cartilage.

**LIP: protein selected for its very high digestibility.

ADDITIVES (KG)**

Nutritional additives: Vitamin A: 19500 IU, Vitamin D3: 700 IU, Iron [3b103]: 33 mg, Iodine [3b201, 3b202]: 3.3 mg, Copper [3b405, 3b406]: 10 mg, Manganese [3b502, 3b504]: 42 mg, Zinc [3b603, 3b605, 3b606]: 135 mg, Selenium [3b801, 3b811, 3b812]: 0.07 mg - Antioxidants.

***Values reflect only levels added to the formula, not those naturally occurring in components of the diet.





URINARY S/O + BLADDER COMFORT



MORSELS IN GRAVY

Available in **box of 12 pouches of 85 g**

RECOMMENDED DAILY INTAKE & FEEDING DURATION

ADULT CAT	CAT'S WEIGHT (kg)	BODY CONDITION			
		NORMAL		OVERWEIGHT	
		grams	pouch	grams	pouch
	2	160	2	125	1 + 1/2
	2.5	185	2	150	2
	3	210	2 + 1/2	170	2
	3.5	235	3	185	2
	4	260	3	205	2 + 1/2
	4.5	280	3 + 1/2	225	2 + 1/2
	5	300	3 + 1/2	240	3
	5.5	325	4	260	3
	6	345	4	275	3
	6.5	365	4 + 1/2	290	3 + 1/2
	7	385	4 + 1/2	305	3 + 1/2
	7.5	405	5	320	4
	8	420	5	335	4
	8.5	440	5	350	4
	9	460	5 + 1/2	365	4 + 1/2
	9.5	475	5 + 1/2	380	4 + 1/2
	10	495	6	395	4 + 1/2

One pouch is equivalent to 18 g of Urinary S/O + Bladder Comfort dry product.

ROYAL CANIN® MULTIFUNCTION URINARY S/O + BLADDER COMFORT morsels in gravy is a complete dietetic feed for cats formulated for the dissolution of struvite stones and the reduction of struvite stone recurrence, and for feline lower urinary tract disease. Urine undersaturating or metastabilising properties for struvite and/or urine acidifying properties.

RECOMMENDATIONS: it is recommended that a veterinarian's opinion be sought before use and before extending the period of use. Feed URINARY S/O + BLADDER COMFORT for 5 to 12 weeks for the dissolution of struvite stones and up to 6 months for the reduction of struvite stone recurrence. Water should be available at all times.

NOTE: In older cats renal function should be checked before recommending a Urinary S/O diet. The use of Urinary S/O Ageing 7+ + Bladder Comfort is recommended in older cats.

[S/O]
[INDEX]

This diet is formulated to promote a healthy urinary environment.



RECOMMENDED FOR CASES OF

- Management of risk factors for Feline Lower Urinary Tract Disease (FLUTD): tendency to be overweight, insufficient water intake and exposure to increased environmental challenges
 - Feline Idiopathic Cystitis (FIC)
 - Struvite uroliths: dissolution and management of recurrence
 - Calcium oxalate uroliths: management of recurrence
 - Stress related urinary conditions
- ### NOT RECOMMENDED FOR CASES OF
- Chronic Kidney Disease (CKD)
 - Heart disease (when sodium restriction is required)
 - Concurrent use of urine-acidifying medications
 - Growth, gestation/lactation



STRUVITE DISSOLUTION

Helps dissolve struvite stones.



HYDRATION SUPPORT

This diet encourages cats' natural drinking behaviour, promoting urine dilution thus limiting urine saturation.



MODERATE CALORIE

A diet with moderate calorie content supports healthy weight maintenance which can help reduce the risk of feline lower urinary tract disease (FLUTD).



BLADDER COMFORT

This diet contains a combination of nutrients to address the leading risk factors of feline lower urinary tract disease including feline idiopathic cystitis (FIC).

KEY VALUES

	Per 1000Kcal	Per 100g as fed
Moisture	1081 g	81.5 g
Protein	106 g	8 g
Fat content	43.77 g	3.3 g
Crude Fibre	15.92 g	1.2 g
Ash	22.55 g	1.7 g
Carbohydrate (NFE)	57.03 g	4.3 g
Dietary Fibre	22.5 g	1.7 g
Omega-6	9.28 g	0.7 g
Omega-3	3.58 g	0.27 g
EPA + DHA	2.12 g	0.16 g
Calcium	1.99 g	0.15 g
Phosphorus	1.72 g	0.13 g
Sodium	3.71 g	0.28 g
Magnesium	0.13 g	0.01 g
Potassium	1.99 g	0.15 g
Copper	3.58 mg	0.27 mg
Chloride	5.31 g	0.4 g
Vitamin A	19629 IU	1480 IU
Vitamin D	292 IU	22 IU
Metabolisable energy*		75.40 Kcal

*Calculated according to NRC 2006 CF.

COMPOSITION

Poultry by-products, pork by-products, salmon, maize flour, pork blood products, powder cellulose, maize starch, minerals, vital wheat gluten, fish oil, hydrolysed milk protein (0.03%), marigold meal, yeasts and part thereof.

ADDITIVES (KG)***

Nutritional additives: Vitamin D3: 50 IU, Taurine: 1 g, Iron [3b103]: 4 mg, Iodine [3b202]: 0.21 mg, Copper [3b405, 3b406]: 1.4 mg, Manganese [3b502, 3b503, 3b504]: 1.2 mg, Zinc [3b603, 3b605, 3b606]: 12 mg - Technological additives: Clnoptilolite of sedimentary origin: 0.4 g.

***Values reflect only levels added to the formula, not those naturally occurring in components of the diet.





RENAL + HYPOALLERGENIC



Available in 2 kg bags

RECOMMENDED DAILY INTAKE & FEEDING DURATION

ADULT CAT	CAT'S WEIGHT (kg)	BODY CONDITION		
		THIN	NORMAL	OVERWEIGHT
		grams	grams	grams
	2	37	31	25
	2.5	43	36	29
	3	50	41	33
	3.5	55	46	37
	4	61	51	41
	4.5	66	55	44
	5	71	59	47
	5.5	76	64	51
	6	81	68	54
	6.5	86	72	57
	7	90	75	60
	7.5	95	79	63
	8	99	83	66
	8.5	104	87	69
	9	108	90	72
	9.5	112	94	75
	10	117	97	78

ROYAL CANIN® RENAL + HYPOALLERGENIC dry is a complete dietetic feed for cats formulated for the support of renal function in the case of chronic renal insufficiency and for the reduction of ingredient and nutrient intolerances. Restricted level of phosphorus. Hydrolysed protein sources and selected carbohydrate sources.

Renal diets are recommended as soon as creatinine goes out of reference range (from IRIS stage 2), before the disease further progresses, and before inappetence develops. When a food allergy or a food intolerance is suspected, the cat's diet should be changed following classical transition. RENAL + HYPOALLERGENIC diet may be continued for the cat's lifetime. Some mineral disturbances (eg. hypokalaemia) may require additional treatment. It is recommended that a veterinarian's opinion be sought before use and before extending the period of use.

It is recommended to divide the daily intake into several meals. Fresh water should be available at all times.



RECOMMENDED FOR CASES OF

Renal disease

- Azotaemic Chronic Kidney Disease (IRIS stages 2 to 4)
- Chronic Kidney Disease with proteinuria (IRIS stages 1 to 4)



Dermatological/GI conditions:

- Adverse Food Reactions (AFR)
- Atopic dermatitis
- Chronic diarrhoea
- Inflammatory Bowel Disease (IBD)
- Exocrine Pancreatic Insufficiency (EPI)

Other:

- Management of calcium oxalate urolith recurrence in cats with impaired renal function
- Management of recurrence of urolithiasis requiring urine alkalinisation: urate & cystine uroliths

NOT RECOMMENDED FOR CASES OF

- Growth, gestation/lactation



RENAL SUPPORT

Formulated with low phosphorus and moderate protein of high quality to support renal function.



ADAPTED ENERGY

Adapted energy content to reduce meal volume and help compensate for a decreased appetite.



HYDROLYSED PROTEIN

Hydrolysed protein with low molecular weight to ensure the food is hypoallergenic.



SKIN BARRIER

Formulated to support the skin's natural protective barrier for optimal skin health.

KEY VALUES

	Per 100Kcal	Per 100g as fed
Moisture	13.41 g	5.5 g
Protein	63.38 g	26 g
Fat content	48.76 g	20 g
Crude Fibre	10 g	4.1 g
Ash	13.9 g	5.7 g
Carbohydrate (NFE)	94.34 g	38.7 g
Dietary Fibre	23.16 g	9.5 g
Omega-6	12.8 g	5.25 g
Omega-3	2.61 g	1.07 g
EPA + DHA	1 g	0.41 g
Calcium	1.46 g	0.6 g
Phosphorus	1 g	0.41 g
Sodium	0.85 g	0.35 g
Chloride	1.95 g	0.8 g
Potassium	2.19 g	0.9 g
Magnesium	0.12 g	0.05 g
Zinc	41.44 mg	17 mg
Vitamin D	195.03	80 IU
Vitamin A	9019.99	3700 IU

Metabolisable energy* 410.2 Kcal

*Calculated according to NRC 2006 CF.

Essential Fatty Acids: Linoleic Acid [5.06%], Arachidonic Acid [0.07%].

COMPOSITION

Rice flour, hydrolysed soya protein isolate, animal fats, rice, chicory pulp, hydrolysed poultry liver, vegetable fibres, minerals, soya oil, fish oil, fructo-oligosaccharides, borage oil, marigold meal.

Carbohydrate sources: rice flour [32.0%], rice [7.7%]. Protein sources: hydrolysed soya protein isolate [23.2%], hydrolysed poultry liver [4.4%].

ADDITIVES (KG)**

Nutritional additives: Vitamin A: 31000 IU, Vitamin D3: 800 IU, Iron [3b103]: 39 mg, Iodine [3b201, 3b202]: 3.5 mg, Copper [3b405, 3b406]: 15 mg, Manganese [3b502, 3b504]: 52 mg, Zinc [3b603, 3b605, 3b606]: 152 mg, Selenium [3b801, 3b811, 3b812]: 0.25 mg - Technological additives: Clinoptilolite of sedimentary origin: 5 g - Antioxidants.

** Values reflect only levels added to the formula, not those naturally occurring in components of the diet.





Available in 1.5 kg bags

RECOMMENDED DAILY INTAKE & FEEDING DURATION

ADULT CAT	IDEAL WEIGHT (kg)	STAGE			
		WEIGHT LOSS		MAINTENANCE AFTER WEIGHT LOSS	
		grams	cups	grams	cups
	2	29	3/8	30	3/8
	2.5	34	3/8	36	4/8
	3	38	4/8	40	4/8
	3.5	43	4/8	45	4/8
	4	47	5/8	50	5/8
	4.5	51	5/8	54	5/8
	5	55	5/8	58	6/8
	5.5	59	6/8	62	6/8
	6	63	6/8	66	6/8
	6.5	66	6/8	70	7/8
	7	70	7/8	74	7/8
	7.5	73	7/8	78	1
	8	77	1	81	1
	8.5	80	1	85	1
	9	84	1	88	1 + 1/8
	9.5	87	1 + 1/8	92	1 + 1/8
	10	90	1 + 1/8	95	1 + 1/8

Weight loss: Individual target weight is determined and then feeding amount established. The recommended daily intake is based on the cat's ideal weight. The starting amount may need to be adjusted during follow-up visits, based on each individual's response and the rate of weight loss. An objective of losing 0.5-2% bodyweight per week is generally achievable over the first 3 months, and of 0.5-1% afterwards.

Maintenance after weight loss: Energy requirements to maintain stable body weight after weight loss are lower than those of cats that have never had obesity. Based on current scientific knowledge, our maintenance after weight loss rations are calculated by increasing the weight loss ration by about 10%.

It is recommended to divide the daily intake into several meals and have fresh water available at all times.

ROYAL CANIN® SATIETY + MOBILITY dry is a complete dietetic feed for cats formulated for the reduction of excessive body weight and for the support of the metabolism of joints in the case of osteoarthritis. Low energy density, high levels of DHA and total omega-3 fatty acids, increased levels of methionine and manganese, and adequate level of vitamin E.

RECOMMENDATIONS: your advice as a veterinarian is necessary before use and before extending the period of use of this diet. Feed this diet until target body weight is achieved and after if required to maintain target body weight. For cats, a transition period when starting the diet is advised. For an efficient weight loss or ideal weight maintenance, the recommended daily energy intake should not be exceeded. For the support of the metabolism of joints in the case of osteoarthritis, feed this diet initially up to 3 months before recheck. The diet may be continued for the cat's lifetime with vet follow-up.

RECOMMENDED FOR CASES OF

Weight management:

- Overweight/obesity: weight loss and maintenance after weight loss
- Stabilised diabetes mellitus with overweight/obesity
- Hyperlipidaemia with overweight/obesity

Joint conditions/mobility:

- Osteoarthritis
- Joints in need of support after orthopaedic surgery or trauma
- Decreased mobility

NOT RECOMMENDED IN CASES OF

- Growth, gestation/lactation
- Underweight
- Conditions where high fibre content is not recommended (eg: chronic constipation/history of constipation in response to a high fibre diet)



EFFECTIVE WEIGHT MANAGEMENT

Provides safe weight loss and helps avoid weight regain. 97% of cats lost weight in 3 months using this nutritional approach.*



BEGGING CONTROL

High level of selected fibres helps keep cats satisfied between meals. Nutritional approach shown to help control begging in 82% of cats during weight loss.*



LOW CALORIE

Overweight can negatively impact mobility. Low calorie content to help achieve and maintain an ideal body weight, thus supporting healthy mobility.



JOINT SUPPORT

With Green Lipped Mussel and a high level of omega-3 fatty acids (EPA+DHA) to support mobility and joints in the case of osteoarthritis.



This diet is formulated to promote a healthy urinary environment.

*Flanagan J et al (2018) An international multi-centre cohort study of weight loss in overweight cats: Differences in outcome in different geographical locations. *PLoS One* :33(7):e0200414.

KEY VALUES

	Per 1000 kcal	Per 100 g as fed
Moisture	17.75 g	5.50 g
Protein	110 g	34 g
Fat content	29.05 g	9 g
Crude fibre	48.10 g	14.90 g
Ash	28.08 g	8.70 g
Carbohydrate (NFE)	90.06 g	27.90 g
Dietary fibre	79.41 g	24.60 g
Omega-6 fatty acids	5.71 g	1.77 g
Omega-3 fatty acids	3.49 g	1.08 g
EPA + DHA	2.68 g	0.83 g
Glucosamine + Chondroitin	323 mg	100 mg
Calcium	4.33 g	1.34 g
Phosphorus	3.39 g	1.05 g
Sodium	1.78 g	0.55 g
Chloride	2.78 g	0.86 g
Magnesium	0.19 g	0.06 g
Vitamin D	323 IU	100 IU
Vitamin E	235.60 IU	730 IU
Taurine	0.61 g	0.19 g
Metabolisable energy**		309.80 kcal

**Calculated according to NRC 2006.

Docosahexaenoic acid (DHA): 0.6% - Methionine: 0.64% - Manganese: 53mg/kg

COMPOSITION

Dehydrated poultry proteins, vegetable fibres, manioc (tapioca), hydrolysed animal proteins, wheat flour, maize gluten, wheat gluten, minerals, chicory pulp, animal fats, algal oil Schizochytrium sp. (source of EPA+DHA), maize flour, psyllium husks and seeds, New Zealand green-lipped mussel powder (GLM) (0.29%), glucosamine from fermentation, marigold meal, fish oil, hydrolysed cartilage (source of chondroitin 0.0005%).

ADDITIVES (KG)***

Nutritional additives: Vitamin A: 25500 IU, Vitamin D3: 1000 IU, Iron (3b103): 29 mg, Iodine (3b201, 3b202): 2.9 mg, Copper (3b405, 3b406): 9 mg, Manganese (3b502, 3b504): 37 mg, Zinc (3b603, 3b605, 3b606): 123 mg, Selenium (3b801, 3b811, 3b812): 0.04 mg - Antioxidants.

*** Values reflect only levels added to the formula, not those naturally occurring in components of the diet.





RENAL + MOBILITY



Available in 2 kg bags

RECOMMENDED DAILY INTAKE & FEEDING DURATION

ADULT CAT	CAT'S WEIGHT (kg)	BODY CONDITION					
		THIN		NORMAL		OVERWEIGHT	
		grams	cups	grams	cups	grams	cups
	2	39	3/8	32	2/8	26	2/8
	2.5	46	3/8	38	3/8	30	2/8
	3	52	4/8	43	3/8	35	3/8
	3.5	58	4/8	48	4/8	39	3/8
	4	64	5/8	53	4/8	42	3/8
	4.5	69	5/8	58	4/8	46	3/8
	5	74	6/8	62	5/8	50	4/8
	5.5	80	6/8	67	5/8	53	4/8
	6	85	6/8	71	5/8	57	4/8
	6.5	90	7/8	75	6/8	60	4/8
	7	95	7/8	79	6/8	63	5/8
	7.5	99	7/8	83	6/8	66	5/8
	8	104	1	87	7/8	69	5/8
	8.5	109	1	91	7/8	72	5/8
	9	113	1	94	7/8	75	6/8
	9.5	118	1 + 1/8	98	7/8	78	6/8
	10	122	1 + 1/8	102	1	81	6/8

It is recommended to divide the daily intake into several meals and have fresh water available at all times.

Renal diets are recommended as soon as creatinine goes out of reference range (from CKD late stage 2), before the disease further progresses, and before inappetence develops. A food transition over 7-10 days is recommended. Cats with CKD IRIS stage 2 that have serum phosphate within IRIS target (< 45 mg/dl) and total calcium over 120 mg/dl should be switched back to a Early Renal diet.¹ Some mineral disturbances (eg. hypokalaemia) may require additional treatment.

ROYAL CANIN® RENAL + MOBILITY dry is a complete dietetic feed for cats formulated for the support of renal function in case of chronic renal insufficiency and for the support of the metabolism of joints in the case of osteoarthritis. High quality proteins and restricted level of phosphorus; high levels of DHA and total omega-3 fatty acids; increased levels of methionine and manganese; and adequate level of vitamin E.

RECOMMENDATIONS: your advice as a veterinarian is necessary before use and before extending the period of use of this diet. For the support of renal function in case of chronic renal insufficiency, feed this diet initially up to 6 months before recheck. For the support of the metabolism of joints in the case of osteoarthritis, feed this diet initially up to 3 months before recheck. The diet may be continued for the cat's lifetime with vet follow-up.

¹ Adapted from IRIS treatment recommendations for CKD in cats (2023).

RECOMMENDED FOR CASES OF

Renal disease:

- Chronic Kidney Disease with proteinuria (IRIS stage 1 to 4)
- Azotaemic Chronic Kidney Disease (IRIS late stage 2 to stage 4)

Joint conditions/mobility:

- Osteoarthritis
- Joints in need of support after an orthopaedic surgery or trauma
- Decreased mobility

Other:

- Management of calcium oxalate urolith recurrence in cats with impaired renal function
- Prevention of recurrence of urolithiasis requiring urine alkalisation: urate & cystine uroliths

NOT RECOMMENDED IN CASES OF

- Growth, gestation/lactation



RENAL SUPPORT

Formulated with low phosphorus and moderate protein of high quality to support renal function, helping to improve the cat's quality of life.



HIGH ENERGY

High energy content to reduce meal volume and help compensate for a decreased appetite.



JOINT SUPPORT

With green-lipped mussel and a high level of omega-3 fatty acids (EPA+DHA) to support mobility and joints in the case of osteoarthritis.



AGEING SUPPORT

Highly digestible formula enriched with vitamins to support healthy ageing.

KEY VALUES

	Per 1000 kcal	Per 100 g as fed
Moisture	14.03 g	5.50 g
Protein	58.66 g	23 g
Fat content	43.36 g	17 g
Crude fibre	11.99 g	4.70 g
Ash	14.79 g	5.80 g
Carbohydrate (NFE)	112 g	44 g
Dietary fibre	27.29 g	10.70 g
Omega-6 fatty acids	8.34 g	3.27 g
Omega-3 fatty acids	3.14 g	1.23 g
EPA + DHA	2.04 g	0.80 g
Glucosamine + Chondroitin	254 mg	99.50 mg
Calcium	1.45 g	0.57 g
Phosphorus	0.74 g	0.29 g
Sodium	1.05 g	0.41 g
Magnesium	0.18 g	0.07 g
Potassium	2.35 g	0.92 g
Vitamin D	204 IU	80 IU
Vitamin E	125 IU	49 IU
Taurine	0.61 g	0.24 g
Metabolisable energy*		392.10 kcal

*Calculated according to NRC 2006.

Essential fatty acids (linoleic acid, arachidonic acid): 3.16% - Docosahexaenoic acid (DHA): 0.55% - Methionine: 0.79% - Manganese: 72 mg/kg

COMPOSITION

Rice, wheat flour, wheat gluten**, animal fats, maize flour, vegetable fibres, maize, soya protein isolate**, minerals, hydrolysed poultry liver, maize gluten*, chicory pulp, algal oil Schizochytrium sp. (source of EPA+DHA), soya oil, dried tomato pulp (source of lycopene), mono- and diglycerides of palmitic and stearic acids esterified with citric acid, psyllium husks and seeds, fructo-oligosaccharides, fish oil, New Zealand green-lipped mussel powder (GLM) (0.3%), glucosamine from fermentation, marigold meal, hydrolysed cartilage (source of chondroitin 0.004%).

**Protein sources: 19.4%.

ADDITIVES (KG)***

Nutritional additives: Vitamin A: 21600 IU, Vitamin D3: 800 IU, Vitamin C: 200 mg, Iron (3b103): 41 mg, Iodine (3b201, 3b202): 4.1 mg, Copper (3b405, 3b406): 13 mg, Manganese (3b502, 3b504): 53 mg, Zinc (3b603, 3b605, 3b606): 149 mg, Selenium (3b801, 3b811, 3b812): 0.07 mg. • Technological additives: Cynoptilolite of sedimentary origin: 10 g. • Sensory additives: Flavours - Antioxidants.





RENAL + MOBILITY



THIN SLICES IN GRAVY

Available in box of 12 pouches of 85 g

RECOMMENDED DAILY INTAKE & FEEDING DURATION

ADULT CAT	CAT'S WEIGHT (kg)	BODY CONDITION					
		THIN		NORMAL		OVERWEIGHT	
		grams	pouches	grams	pouches	grams	pouches
2	120	1 + 1/2	1	100	1	80	1
2.5	140	1 + 1/2	1	120	1 + 1/2	95	1
3	160	2	1	135	1 + 1/2	110	1 + 1/2
3.5	180	2	1	150	2	120	1 + 1/2
4	200	2 + 1/2	1	165	2	130	1 + 1/2
4.5	215	2 + 1/2	1	180	2	145	1 + 1/2
5	230	2 + 1/2	1	195	2 + 1/2	155	2
5.5	250	3	1	205	2 + 1/2	165	2
6	265	3	1	220	2 + 1/2	175	2
6.5	280	3 + 1/2	1	235	3	185	2
7	295	3 + 1/2	1	245	3	195	2 + 1/2
7.5	310	3 + 1/2	1	260	3	205	2 + 1/2
8	325	4	1	270	3	215	2 + 1/2
8.5	340	4	1	285	3 + 1/2	225	2 + 1/2
9	355	4	1	295	3 + 1/2	235	3
9.5	365	4 + 1/2	1	305	3 + 1/2	245	3
10	380	4 + 1/2	1	315	3 + 1/2	255	3

It is recommended to divide the daily intake into several meals and have fresh water available at all times.

Renal diets are recommended as soon as creatinine goes out of reference range (from CKD late stage 2), before the disease further progresses, and before inappetence develops. A food transition over 7-10 days is recommended. Cats with CKD IRIS stage 2 that have serum phosphate within IRIS target (≤ 4.5 mg/dL) and total calcium over 120 mg/dL should be switched back to a Early Renal diet. Some mineral disturbances (eg hypocalcaemia) may require additional treatment.

ROYAL CANIN® RENAL + MOBILITY thin slices in gravy is a complete dietetic feed for cats formulated for the support of renal function in case of chronic renal insufficiency and for the support of the metabolism of joints in the case of osteoarthritis. High quality proteins and restricted level of phosphorus, high levels of DHA and total omega-3 fatty acids, increased levels of methionine and manganese, and adequate level of vitamin E.

RECOMMENDATIONS: your advice as a veterinarian is necessary before use and before extending the period of use of this diet. For the support of renal function in case of chronic renal insufficiency, feed this diet initially up to 6 months before recheck. For the support of the metabolism of joints in the case of osteoarthritis, feed this diet initially up to 3 months before recheck. The diet may be continued for the cat's lifetime with vet follow-up.

RECOMMENDED FOR CASES OF

Renal disease:

- Chronic Kidney Disease with proteinuria (IRIS stage 1 to 4)
- Azotaemic Chronic Kidney Disease (IRIS late stage 2 to stage 4)

Joint conditions/mobility:

- Osteoarthritis
- Joints in need of support after orthopaedic surgery or trauma
- Decreased mobility

Other:

- Management of calcium oxalate urolith recurrence in cats with impaired renal function
- Prevention of recurrence of urolithiasis requiring urine alkalinisation: urate & cystine uroliths

NOT RECOMMENDED IN CASES OF

- Growth, gestation/lactation



RENAL SUPPORT

Formulated with low phosphorus and moderate protein of high quality to support renal function, helping to improve the cat's quality of life.



HIGH ENERGY

High energy content to reduce meal volume and help compensate for a decreased appetite.



JOINT SUPPORT

High level of omega-3 fatty acids (EPA+DHA) to support mobility and joints in the case of osteoarthritis.



AGEING SUPPORT

Highly digestible formula enriched with vitamins to support healthy ageing.

KEY VALUES

	Per 1000 kcal	Per 100 g as fed
Moisture	658.12 g	77 g
Protein	65.81 g	7.70 g
Fat content	67.52 g	7.90 g
Crude fibre	3.42 g	0.40 g
Ash	12.82 g	1.50 g
Carbohydrate (NFE)	47.01 g	5.50 g
Dietary fibre	6.84 g	0.80 g
Omega-6 fatty acids	20.51 g	2.40 g
Omega-3 fatty acids	3.59 g	0.42 g
EPA + DHA	2.31 g	0.27 g
Glucosamine + Chondroitin	270.09 mg	31.60 mg
Calcium	1.28 g	0.15 g
Phosphorus	0.77 g	0.09 g
Sodium	1.03 g	0.12 g
Magnesium	0.17 g	0.02 g
Potassium	1.88 g	0.22 g
Vitamin D	376.10 IU	44 IU
Vitamin E	132.50 IU	15.50 IU
Taurine	1.28 g	0.15 g
Metabolisable energy*		117 kcal

*Calculated according to NRC 2006.

Essential fatty acids (linoleic acid, arachidonic acid): 2.25% - Docosahexaenoic acid (DHA): 0.17% - Methionine: 0.15% - Manganese: 4.4 mg/kg

COMPOSITION

Poultry by-products**, wheat flour, pork blood products**, sunflower oil refined, maize flour, modified starch, minerals, algal oil Schizochytrium sp. (source of EPA+DHA), fish oil, powder cellulose, dried beet pulp, dried tomato pulp, fructo-oligosaccharides, glucosamine from fermentation, marigold meal, hydrolysed cartilage (source of chondroitin(0.001%)).

**Protein sources: 28.2%

ADDITIVES (KG)***

Nutritional additives: Vitamin D3: 390 IU, Vitamin C: 62 mg, Taurine: 1.5 g, Iron (3b103): 6 mg, Iodine (3b202): 0.62 mg, Copper (3b405, 3b406): 4.1 mg, Manganese (3b502, 3b503, 3b504): 1.9 mg, Zinc (3b603, 3b605, 3b606): 19 mg - Technological additives: Clnoptilolite of sedimentary origin: 2 g - Sensory additives: Flavourings.

*** Values reflect only levels added to the formula, not those naturally occurring in components of the diet.

¹ Adapted from IRIS treatment recommendations for CKD in cats (2012).





CARDIAC



Available in 1.5 kg bags

RECOMMENDED DAILY INTAKE & FEEDING DURATION

ADULT CAT	CAT'S WEIGHT (kg)	BODY CONDITION		
		THIN	NORMAL	OVERWEIGHT
		grams	grams	grams
	2	41	34	27
	2.5	48	40	32
	3	54	45	36
	3.5	60	50	40
	4	66	55	44
	4.5	72	60	48
	5	78	65	52
	5.5	83	70	56
	6	89	74	59
	6.5	94	78	63
	7	99	83	66
	7.5	104	87	69
	8	109	91	73
	8.5	114	95	76
	9	118	99	79
	9.5	123	103	82
	10	127	106	85

ROYAL CANIN® CARDIAC dry is a complete dietetic feed for cats formulated for the support of heart function in the case of chronic cardiac insufficiency. Restricted level of sodium.

RECOMMENDATIONS: It is recommended that advice from a veterinarian be sought before use and before extending the period of use. Feed CARDIAC dry initially up to 6 months.



This diet is formulated to promote a healthy urinary environment.

RECOMMENDED FOR CASES OF

- Hypertrophic cardiomyopathy
- Congestive heart failure

NOT RECOMMENDED IN CASE OF

- Chronic Kidney Disease (CKD)
- Growth, gestation/lactation



SUPPORTS HEART FUNCTION

Unique formula enriched with vitamin E, vitamin C, taurine and omega-3 fatty acids (EPA+DHA) to support heart function.



REDUCES HEART WORKLOAD

Restricted level of sodium to help support heart function in the case of chronic cardiac insufficiency.



URINARY HEALTH SUPPORT

Adapted to support the needs of neutered cats with urinary sensitivities.

KEY VALUES

	Per 1000Kcal	Per 100g as fed
Moisture	14.67 g	5.5 g
Protein	121.87 g	45.7 g
Fat content	34.67 g	13 g
Crude Fibre	17.33 g	6.5 g
Ash	18.40 g	6.9 g
Carbohydrate (NFE)	59.73 g	22.4 g
Dietary Fibre	41.6 g	15.6 g
Omega-6	7.49 g	2.81 g
Omega-3	1.55 g	0.58 g
EPA + DHA	0.72 g	0.27 g
Calcium	3.2 g	1.2 g
Phosphorus	2.4 g	0.9 g
Magnesium	0.19 g	0.07 g
Sodium	0.67 g	0.25 g
Potassium	1.87 g	0.7 g
Chloride	2.29 g	0.86 g
Taurine	0.59 g	0.22 g
Vitamin E	116.53 mg	43.7 mg
Vitamin C	47.47 mg	17.8 mg
Metabolisable energy*		375 Kcal*

Calculated according to NRC 2006 CF.

COMPOSITION

Dehydrated poultry protein, barley, wheat gluten**, maize gluten, vegetable fibres, animal fats, hydrolysed animal proteins, chicory pulp, yeasts products, minerals, psyllium husks and seeds, fish oil, mono- and diglycerides of palmitic and stearic acids esterified with citric acid, soya oil, marigold meal.

**LIP (Low Indigestible Protein): protein selected because of its very high digestibility.

ADDITIVES (KG)**

Nutritional additives: Vitamin A: 18554 IU, Vitamin D3: 964 IU, Vitamin E: 437 mg, Vitamin C: 178 mg, Taurine: 2.2 g, Iron [3b103]: 19 mg, Iodine [3b201, 3b202]: 1.9 mg, Copper [3b405, 3b406]: 5.9 mg, Manganese [3b502, 3b504]: 25 mg, Zinc [3b603, 3b605, 3b606]: 113 mg, Selenium [3b801, 3b811, 3b812]: 0.05 mg, Zootechnical Additives: Ammonium chloride: 5 g - Antioxidants.

*** Values reflect only levels added to the formula, not those naturally occurring in components of the diet.





GASTROINTESTINAL

Hydrolysed protein



This diet supports a healthy urinary environment.

S/O INDEX

Available in 2kg, 4kg

RECOMMENDED DAILY INTAKE

ADULT CAT	CAT'S WEIGHT (kg)	WEIGHT CONDITION		
		UNDERWEIGHT	IDEAL WEIGHT	OVERWEIGHT
		grams	grams	grams
	2	38	31	25
	2.5	44	37	29
	3	50	42	33
	3.5	56	47	37
	4	62	51	41
	4.5	67	56	45
	5	72	60	48
	5.5	77	65	52
	6	82	69	55
	6.5	87	73	58
	7	92	77	61
	7.5	96	80	64
	8	101	84	67

It is recommended to divide the daily intake into several small meals throughout the day and have fresh water available at all times.

For acute cases, feeding a GI diet at least 2-3 weeks (continuing after clinical signs have resolved) helps support optimal pet recovery and regeneration of the intestinal villi. If signs have not resolved in 5-7 days, consider additional diagnostics such as blood tests, imaging and infectious disease testing.

For chronic enteropathies, nutritional management is key. Successive 2-week dietary trials after a 7-10 day transition should be conducted until the most effective diet is identified. After a 2-week trial is completed: if full remission achieved, continue diet long term, in case of partial remission, continue trial for 2 more weeks and if there is no improvement, try a different GI diet. If no diet leads to clinical remission, the most effective diet at reducing signs should be continued long term alongside appropriate medications.

ROYAL CANIN® GASTROINTESTINAL HYDROLYSED PROTEIN dry is a complete dietetic feed for cats formulated for the reduction of ingredient and nutrient intolerances, for the compensation for maldigestion and for the reduction of intestinal absorptive disorders. Hydrolysed protein sources and selected carbohydrate sources. Highly digestible feed with increased sodium and potassium.

RECOMMENDATIONS: your advice as a veterinarian is necessary before use and before extending the period of use of this diet. For the reduction of ingredient and nutrient intolerances, feed this diet 3 to 8 weeks before recheck and up to one year before reassessing nutritional recommendation if signs of intolerance disappear. For the compensation for maldigestion and for the reduction of intestinal absorptive disorders, feed this diet initially up to 12 weeks before recheck and lifetime in the case of chronic pancreatic insufficiency. The diet may be continued for the cat's lifetime with vet follow-up.

PROVIDES DIGESTIVE SUPPORT WHILE MINIMISING THE RISK OF ADVERSE FOOD REACTIONS.

RECOMMENDED FOR CASES OF

- Chronic Inflammatory Enteropathy (CIE) - formerly called Inflammatory Bowel Disease (IBD)
- Acute and chronic diarrhoea
- Maldigestion/malabsorption including Exocrine Pancreatic Insufficiency (EPI)
- Adverse Food Reactions (AFR)
- Impaired appetite, feeding post surgery
- Acute hepatopathy/hepatic lipidosis without hepatic encephalopathy

NOT RECOMMENDED IN CASE OF

- Hyperlipidemia¹
- Hepatic encephalopathy²
- Gestation or lactation³

¹ In these cases the recommended diet is Gastrointestinal Moderate Calorie - ² In these cases the recommended diet is Hepatic - ³ For kittens or gestating/lactating queens the most appropriate is Gastrointestinal Kitten

For kittens, can be used under vet supervision, with close follow-up and monitoring of growth.



DIGESTIVE SUPPORT

A highly digestible formula with balanced fibres to support a healthy digestion and transit.



MICROBIOME SUPPORT

Formulated with selected prebiotics to support a healthy gut and intestinal microbiome.



HIGH ENERGY

High energy content to reduce meal volume and help compensate for a decreased appetite.



HYDROLYSED PROTEIN

Highly digestible hydrolysed protein to ensure the food is hypoallergenic.

KEY VALUES

	Per 1000 kcal	Per 100 g as fed
Moisture	13.61 g	5.5 g
Protein	76.69 g	31 g
Fat content	49.48 g	20 g
Crude Fibre	11.38 g	4.6 g
Ash	18.8 g	7.6 g
Carbohydrate (NFE)	77.44 g	31.3 g
Total Dietary Fiber (TDF)	25.98 g	10.5 g
Omega-6 fatty acids	9.20 g	3.72 g
Omega-3 fatty acids	2.28 g	0.92 g
EPA + DHA	1.01 g	0.41 g
Calcium	1.73 g	0.70 g
Phosphorus	1.48 g	0.60 g
Potassium	2.47 g	1 g
Sodium	1.61 g	0.65 g
Chloride	2.85 g	1.15 g
Magnesium	0.17 g	0.07 g
Copper	5.12 mg	2.07 mg
Zinc	42.06 mg	17mg
Metabolisable energy*		404.2 kcal

*Calculated according to NRC 2006 CF - Essential fatty acids (linoleic acid, arachidonic acid): 3.68%

COMPOSITION

Hydrolysed soya protein isolate*, rice flour**, animal fats***, vegetable fibres, animal protein hydrolysate***, minerals, broken rice**, beet pulp, fish oil***, fructo-oligosaccharides, psyllium husks, yeast components (source of manno-oligosaccharides 0.05%), marigold meal.

*Protein source: 29.1%. **Carbohydrate sources: 31.1%. ***Highly digestible ingredients: 32.6%.

ADDITIVES (KG)**

Nutritional additives: Vitamin A: 34300 IU, Vitamin D3: 800 IU, Vitamin E: 809 mg, Vitamin C: 249 mg, Iron (3b103): 39 mg, Iodine (3b201, 3b202): 3.5 mg, Copper (3b405): 14.4 mg, Manganese (3b502, 3b505): 51 mg, Zinc (3b603, 3b605, 3b612): 151 mg, Selenium (3b601, 3b611, 3b612): 0.24 mg - Technological additives: Cimplotilolite of sedimentary origin: 10 g - Antioxidants.

**Values reflect only levels added to the formula, not those naturally occurring in components of the diet.



VET SERVICES

Scan to access the Vet Services Feeding Amount Calculator for a precise, individualized feeding recommendation.



GLYCOADVANCED



GLYCOADVANCED



Available in 1.5 and 3.5 kg bags

RECOMMENDED DAILY INTAKE & FEEDING DURATION

ADULT CAT	IDEAL WEIGHT / TARGET* WEIGHT (KG)	HEALTHY WEIGHT MAINTENANCE	STARTING WEIGHT LOSS	MAINTENANCE AFTER WEIGHT LOSS
		grams	grams	grams
	3	51	35	40
	3.5	57	39	44
	4	63	43	49
	4.5	68	47	53
	5	74	50	57
	5.5	79	54	61
	6	84	57	65
	6.5	89	61	69
	7	93	64	72
	7.5	98	67	76
	8	103	70	79
	8.5	107	73	83
	9	112	76	86
	9.5	116	79	90
	10	120	82	93

*Cats with BCS > 4/9 should initially be fed for their current weight. Once clinical signs are stable and fair to good glycaemic control has been achieved, caloric restriction can be undertaken (feeding volume for caloric restriction should be based on the cat's ideal weight). For cats undergoing weight reduction, we recommend a follow-up consultation 2-4 weeks after starting the diet to assess their individual response. **NOTE: The use of this food may affect the cat's sensitivity to insulin. In insulin treated cats, insulin dosage may require adjustment in response to dietary change; it is highly recommended that blood glucose and insulin responses are monitored when starting or changing the diet, including changes in feeding amounts. In the management of diabetes mellitus, it is recommended to keep the type of food and daily rationing as consistent as possible. Divide the daily ration into several meals and have fresh water available at all times.**

ROYAL CANIN® GLYCOADVANCED dry is a complete dietetic feed for cats formulated for the reduction of excessive body weight and for the regulation of glucose supply (diabetes mellitus). Low energy density and low level of mono- and disaccharides.

RECOMMENDATIONS: your advice as a veterinarian is necessary before use and before extending the period of use of this diet. Feed this diet until target body weight is achieved and after, if required, to maintain target body weight. For cats, a transition period when starting the diet is advised. For efficient weight loss or ideal weight maintenance, the recommended daily energy intake should not be exceeded. For the regulation of glucose supply, feed this diet initially for up to 6 months. The diet may be continued for the cat's lifetime with veterinary follow-up.



RECOMMENDED FOR CASES OF

- Diabetes mellitus and Body Condition Score (BCS) > 5/9

NOT RECOMMENDED IN CASE OF

- Underweight (BCS < 5/9)
- Growth, gestation/lactation

Proven in a 12-week clinical study with a veterinarian-approved diabetic management programme:



TO SUPPORT GLUCOMODULATION

To support the regulation of blood glucose levels.



TO SUPPORT GLUCOSE METABOLISM

To help resume physiological glucose management in a majority of diabetic cats undergoing a weight loss programme.



TO PROMOTE HEALTHY WEIGHT

To support complementary beneficial effects in both weight and glucose management in overweight diabetic cats.



[S/O]
INDEX

This diet is formulated to promote a healthy urinary environment.

KEY VALUES

	Per 1000 kcal	Per 100g as fed
Moisture	16.12 g	5.50 g
Protein	138 g	47 g
Fat content	26.38 g	9 g
Crude fibre	28.14 g	9.60 g
Crude ash	23.15 g	7.90 g
Carbohydrate (NFE)	61.55 g	21 g
Total dietary fibre	58.32 g	19.90 g
Omega-6 fatty acids	5.74 g	1.96 g
Omega-3 fatty acids	1.47 g	0.50 g
EPA + DHA	0.88 g	0.30 g
Calcium	2.81 g	0.96 g
Phosphorus	2.49 g	0.85 g
Potassium	3.66 g	1.25 g
Sodium	1.82 g	0.62 g
Chloride	4.19 g	1.43 g
Magnesium	0.18 g	0.06 g
Copper	4.98 mg	1.70 mg
Taurine	0.59 g	0.20 g
L-carnitine	70.34 mg	24 mg
Metabolisable energy**		341.20 Kcal

**Calculated according to NRC 2006 CF.

COMPOSITION

Dehydrated poultry proteins, wheat gluten, vegetable fibres, soya protein isolate, barley***, hydrolysed animal proteins, maize gluten, manioc (tapioca)***, psyllium husks and seeds, minerals, animal fats, chicory pulp, yeast products, algal oil Schizochytrium sp., fish oil, marigold meal.

***Carbohydrate sources: 11.1%.

ADDITIVES (KG)****

Nutritional additives: Vitamin A: 24800 IU, Vitamin D3: 1000 IU, Iron [3b103]: 25 mg, Iodine [3b201, 3b202]: 2.5 mg, Copper [3b405, 3b406]: 7.8 mg, Manganese [3b502, 3b504]: 33 mg, Zinc [3b603, 3b605, 3b606]: 100 mg, Selenium [3b801, 3b811, 3b812]: 0.03 mg - Antioxidants.

**** Values reflect only levels added to the formula, not those naturally occurring in components of the diet.



GLYCOADVANCED



MORSELS IN GRAVY

Available in a box of 12 pouches of 85 g

RECOMMENDED DAILY INTAKE & FEEDING DURATION

	IDEAL WEIGHT / TARGET* WEIGHT (KG)	HEALTHY WEIGHT MAINTENANCE		STARTING WEIGHT LOSS		MAINTENANCE AFTER WEIGHT LOSS	
		grams	pouch	grams	pouch	grams	pouch
ADULT CAT	3	275	3	190	2	215	2 + 1/2
	3.5	310	3 + 1/2	210	2 + 1/2	240	3
	4	340	4	230	2 + 1/2	265	3
	4.5	370	4 + 1/2	250	3	285	3 + 1/2
	5	400	4 + 1/2	270	3	310	3 + 1/2
	5.5	425	5	290	3 + 1/2	330	4
	6	455	5 + 1/2	310	3 + 1/2	350	4
	6.5	480	5 + 1/2	330	4	370	4 + 1/2
	7	505	6	345	4	390	4 + 1/2
	7.5	530	6	365	4 + 1/2	410	5
	8	555	6 + 1/2	380	4 + 1/2	430	5
	8.5	580	7	395	4 + 1/2	450	5 + 1/2
	9	605	7	415	5	470	5 + 1/2
	9.5	630	7 + 1/2	430	5	485	5 + 1/2
	10	650	7 + 1/2	445	5	505	6

*Cats with BCS $\geq 6/9$ should initially be fed for their current weight. Once clinical signs are stable and fair to good glycaemic control has been achieved, caloric restriction can be undertaken (feeding volume for caloric restriction should be based on the cat's ideal weight). For cats undergoing weight reduction, we recommend a follow-up consultation 2-4 weeks after starting the diet to assess their individual response.

NOTE: The use of this food may affect the cat's sensitivity to insulin. In insulin treated cats, insulin dosage may require adjustment in response to dietary change; it is highly recommended that blood glucose and insulin responses are monitored when starting or changing the diet, including changes in feeding amounts. In the management of diabetes mellitus, it is recommended to keep the type of food and daily rationing as consistent as possible. Divide the daily ration into several meals and have fresh water available at all times.

ROYAL CANIN® GLYCOADVANCED morsels in gravy is a complete dietetic feed for cats formulated for the reduction of excessive body weight and for the regulation of glucose supply (diabetes mellitus). Low energy density and low level of mono- and disaccharides.

RECOMMENDATIONS: your advice as a veterinarian is necessary before use and before extending the period of use of this diet. Feed this diet until target body weight is achieved and after, if required, to maintain target body weight. For cats, a transition period when starting the diet is advised. For efficient weight loss or ideal weight maintenance, the recommended daily energy intake should not be exceeded. For the regulation of glucose supply, feed this diet initially for up to 6 months. The diet may be continued for the cat's lifetime with veterinary follow-up.



RECOMMENDED FOR CASES OF

- Diabetes mellitus and Body Condition Score (BCS) $\geq 5/9$

NOT RECOMMENDED IN CASE OF

- Underweight (BCS $< 5/9$)
- Growth, gestation/lactation

Proven in a 12-week clinical study with a veterinarian-approved diabetic management programme:



TO SUPPORT GLUCOMODULATION

To support the regulation of blood glucose levels.



TO SUPPORT GLUCOSE METABOLISM

To help resume physiological glucose management in a majority of diabetic cats undergoing a weight loss programme.



TO PROMOTE HEALTHY WEIGHT

To support complementary beneficial effects in both weight and glucose management in overweight diabetic cats.



[S/O® INDEX] This diet is formulated to promote a healthy urinary environment.

KEY VALUES

	Per 1000 kcal	Per 100g as fed
Moisture	1446 g	84 g
Protein	170 g	9.90 g
Fat content	27.54 g	1.60 g
Crude fibre	30.98 g	1.80 g
Crude ash	24.10 g	1.40 g
Carbohydrate [NFE]	22.38 g	1.30 g
Total dietary fibre	43.03 g	2.50 g
Omega-6 fatty acids	6.88 g	0.40 g
Omega-3 fatty acids	1.38 g	0.08 g
EPA + DHA	1.03 g	0.06 g
Calcium	2.07 g	0.12 g
Phosphorus	1.72 g	0.10 g
Potassium	3.44 g	0.20 g
Sodium	2.41 g	0.14 g
Chloride	4.99 g	0.29 g
Magnesium	0.33 g	0.02 g
Copper	7.57 mg	0.44 mg
Taurine	1.20 g	0.07 g
L-carnitine	74.01 mg	4.30 mg
Metabolisable energy**		58.10 Kcal

**Calculated according to NRC 2006 CF.


COMPOSITION

Poultry by-products, pork by-products, pork blood products, powder cellulose, hydrolysed pork collagen, minerals, dried beet pulp, algal oil Schizochytrium sp., fish oil, psyllium husks and seeds, marigold meal.

ADDITIVES (KG)***

Nutritional additives: Vitamin D3: 50 IU, Taurine: 0.7 g, Iron (3b103): 5 mg, Iodine (3b202): 0.35 mg, Copper (3b405, 3b406): 2.3 mg, Manganese (3b502, 3b503, 3b504): 1.5 mg, Zinc (3b603, 3b605, 3b606): 15 mg - Technological additives: Clinoptilolite of sedimentary origin: 1 g - Sensory additives: Flavours.

*** Values reflect only levels added to the formula, not those naturally occurring in components of the diet.



URINARY S/O + SATIETY



Available in 3.5, 9 and 15 kg bags

RECOMMENDED DAILY INTAKE & FEEDING DURATION

ADULT DOG	IDEAL DOG'S WEIGHT (kg)	BODY CONDITION		
		STARTING WEIGHT LOSS	AFTER 4 WEEKS	MAINTENANCE AFTER WEIGHT LOSS
		grams	grams	grams
	2	44 g	38 g	42 g
	4	74 g	64 g	70 g
	6	101 g	86 g	95 g
	8	125 g	107 g	118 g
	10	148 g	127 g	139 g
	15	201 g	172 g	189 g
	20	249 g	213 g	235 g
	25	294 g	252 g	277 g
	30	337 g	289 g	318 g
	35	379 g	324 g	357 g
	40	418 g	359 g	394 g
	45	457 g	392 g	431 g
	50	495 g	424 g	466 g
	55	531 g	455 g	501 g
	60	567 g	486 g	535 g
	70	637 g	546 g	600 g
	80	704 g	603 g	663 g

ROYAL CANIN® URINARY S/O + SATIETY dry is a complete dietetic feed for dogs, formulated for the dissolution of struvite stones, the reduction of struvite stone recurrence and for the reduction of excessive body weight. Urine undersaturating or metastabilising properties for struvite and/or Urine acidifying properties.

RECOMMENDATIONS: It is recommended that advice from a veterinarian be sought before use and before extending the period of use. Feed URINARY S/O + SATIETY dry for 5 to 12 weeks for the dissolution of struvite stones, initially up to 6 months for the reduction of struvite stone recurrence, as well as until the target body weight is achieved and after if required to maintain target body weight. For an efficient weight loss or ideal weight maintenance, the recommended daily energy intake should not be exceeded.



This diet is formulated to promote a healthy urinary environment.



RECOMMENDED FOR CASES OF

Urinary disorders:

- Struvite uroliths: dissolution and management of recurrence
- Calcium oxalate uroliths: management of recurrence
- Bacterial cystitis: management of secondary struvite crystalluria (in association with adequate antibiotic therapy)

Weight management:

- Overweight/obesity
- Stabilised Diabetes Mellitus with overweight/obesity
- Hyperlipidaemia with overweight/obesity

NOT RECOMMENDED FOR CASES OF

- Chronic Kidney Disease [CKD]
- Cardiac disease (when sodium restriction is recommended) /Heart failure
- Concurrent use of urine-acidifying medications
- Conditions where high fibre content is not recommended (eg: chronic constipation, severe arthritis making defaecation difficult)
- Growth, gestation/lactation



URINE DILUTION

Urine dilution makes the urine less liable to form crystals.



LOW RSS

Helps lower the concentration of ions contributing to crystal formation.



HIGH PROTEIN

A high protein content helps maintain muscle mass throughout the weight loss period.



SATIETOGENIC EFFECT

Formulated with a special blend of fibres to promote satiety and reduce the spontaneous consumption of food.

KEY VALUES

	Per 1000Kcal	Per 100g as fed
Moisture	35.7 g	9.5 g
Protein	113 g	30 g
Fat content	33.82 g	9 g
Crude Fibre	61.26 g	16.3 g
Ash	27.43 g	7.3 g
Carbohydrate [NFE]	104.85 g	27.9 g
Dietary Fibre	104 g	27.6 g
Omega-6	7.48 g	1.99 g
Omega-3	2.03 g	0.54 g
EPA + DHA	1.05 g	0.28 g
Calcium	3.38 g	0.9 g
Phosphorus	2.63 g	0.7 g
Potassium	3.01 g	0.8 g
Sodium	3.76 g	1 g
Chloride	6.01 g	1.6 g
Magnesium	0.23 g	0.06 g
Copper	6.39 mg	1.7 mg
Metabolisable energy*		266.1 Kcal

*Calculated according to NRC 2006 TDF.
Sulphur 0.7%.

COMPOSITION

Dehydrated poultry proteins, cellulose powder, wheat gluten**, manioc (tapioca), vegetable fibres, hydrolysed animal proteins, wheat, chicory pulp, maize, animal fats, minerals, maize gluten, fish oil, psyllium husks and seeds, fructo-oligosaccharides**, soya oil, sodium butyrate, marigold meal, glucosamine from fermentation, hydrolysed cartilage [source of chondroitin].

**LIP [Low Indigestible Protein]: protein selected because of its very high digestibility.

ADDITIONS (KG)***

Nutritional additives: Vitamin A: 19900 IU, Vitamin D3: 1000 IU, Iron [3b103]: 40 mg, Iodine [3b201, 3b202]: 4 mg, Copper [3b405, 3b406]: 12 mg, Manganese [3b502, 3b504]: 52 mg, Zinc [3b603, 3b605, 3b606]: 137 mg, Selenium [3b801, 3b811, 3b812]: 0.07 mg - Preservatives - Antioxidants.

*** Values reflect only levels added to the formula, not those naturally occurring in components of the diet.



Available in 1 and 3 kg bags

RECOMMENDED DAILY INTAKE & FEEDING DURATION

	IDEAL WEIGHT (kg)	STAGE			
		WEIGHT LOSS		MAINTENANCE AFTER WEIGHT LOSS	
		grams	cups	grams	cups
ADULT DOG	2	38	3/8	42	4/8
	2.5	45	4/8	49	4/8
	3	51	4/8	57	5/8
	3.5	58	5/8	63	5/8
	4	64	5/8	70	6/8
	4.5	70	6/8	77	7/8
	5	75	6/8	83	7/8
	5.5	81	7/8	89	1
	6	86	7/8	95	1
	6.5	92	1	101	1 + 1/8
	7	97	1	107	1 + 1/8
	7.5	102	1 + 1/8	112	1 + 2/8
	8	107	1 + 1/8	118	1 + 2/8
	8.5	112	1 + 2/8	124	1 + 3/8
	9	117	1 + 2/8	129	1 + 3/8
	9.5	122	1 + 2/8	134	1 + 3/8
	10	127	1 + 3/8	140	1 + 4/8

Weight loss: Individual target weight is determined and then feeding amount established. The recommended daily intake is based on the dog's ideal weight. The starting ration may need to be adjusted during follow-up visits, based on each individual's response and the rate of weight loss. An objective of losing 1-3% bodyweight per week is generally achievable over the first 3 months, and of 0.5-2% afterwards. Maintenance after weight loss: Energy requirements to maintain stable body weight after weight loss are lower than those of dogs that have never had obesity. Based on current scientific knowledge, our maintenance after weight loss rations are calculated by increasing the weight loss ration by about 10%.

It is recommended to divide the daily intake into at least 2 meals and have fresh water available at all times.

ROYAL CANIN® URINARY S/O + SATIETY SMALL DOGS dry is a complete dietetic feed for dogs formulated for the dissolution of struvite stones, the reduction of struvite stone recurrence and for the reduction of excessive body weight. Urine undersaturating or metabolising properties for struvite and/or urine acidifying properties. RECOMMENDATIONS: your advice as a veterinarian is necessary before use and before extending the period of use of this diet. Feed this diet 5 to 12 weeks for the dissolution of struvite stones and initially up to 6 months before recheck for the reduction of struvite stone recurrence. Feed until target body weight is achieved and after if required to maintain target body weight. For an efficient weight loss or ideal weight maintenance, the recommended daily energy intake should not be exceeded.

The diet may be continued for the dog's lifetime with vet follow-up.



RECOMMENDED FOR CASES OF

Urinary disorders:

- Struvite uroliths: dissolution and management of recurrence
- Calcium oxalate uroliths: management of recurrence
- Bacterial cystitis: management of secondary struvite crystalluria (in association with adequate antibiotic therapy)

Weight management:

- Overweight/obesity: weight loss and maintenance after weight loss
- Stabilised diabetes mellitus with concomitant overweight/obesity
- Hyperlipidaemia with concomitant overweight/obesity

NOT RECOMMENDED FOR CASES OF

- Growth, gestation or lactation
- Azotaemic Chronic Kidney Disease (IRIS stage 2 to 4)
- Concurrent use of urine-acidifying drugs
- Congestive Heart Failure/Cardiac Disease (when sodium restriction is sought)
- Conditions where high fibre content is not recommended (eg: chronic constipation, severe arthritis making defecation difficult)



URINE DILUTION

Urine dilution makes the urine less liable to form crystals.



LOW RSS

Helps lower the concentration of ions contributing to crystal formation.



HIGH PROTEIN

A high protein content helps maintain muscle mass throughout the weight loss period.



SATIETOGENIC EFFECT

Formulated with a special blend of fibres to promote satiety and reduce the spontaneous consumption of food.



SMALL KIBBLE

Carefully calibrated kibble size to fit small jaws.

[S/O] This diet is formulated to create an environment unfavourable to the formation of struvite and calcium oxalate.

KEY VALUES

	Per 1000 kcal	Per 100g as fed
Moisture	37.05 g	9.50 g
Protein	117 g	30 g
Fat content	37.05 g	9.50 g
Crude Fibre	63.57 g	16.30 g
Ash	28.47 g	7.30 g
Carbohydrate (NFE)	107 g	27.40 g
Dietary Fibre	108 g	27.60 g
Omega-6 fatty acids	7.68 g	1.97 g
Omega-3 fatty acids	2.11 g	0.54 g
EPA + DHA	1.09 g	0.28 g
Calcium	3.39 g	0.87 g
Phosphorus	2.73 g	0.70 g
Potassium	3.12 g	0.80 g
Sodium	3.63 g	0.93 g
Chloride	6.05 g	1.55 g
Magnesium	0.23 g	0.06 g
Copper	6.63 mg	1.70 mg
Metabolisable energy*		256.40 kcal

*Calculated according to NRC 2006 CF.

Sulphur 0.64%

COMPOSITION

Dehydrated poultry proteins, cellulose powder, wheat gluten**, manioc (topical), vegetable fibres, hydrolysed animal proteins, wheat, chicory pulp, maize, animal fats, minerals, maize gluten, maize flour, fish oil, psyllium husks and seeds, soya oil, sodium butyrate, fructooligosaccharides, marigold meal, glucosamine from fermentation, hydrolysed cartilage (source of chondroitin).

**L.I.P.: protein selected for its very high digestibility.

ADDITIVES (KG)***

Nutritional additives: Vitamin A: 19900 IU, Vitamin D3: 1000 IU, Iron [3b103]: 40 mg, Iodine [3b201, 3b202]: 4 mg, Copper [3b405, 3b406]: 12 mg, Manganese [3b502, 3b504]: 52 mg, Zinc [3b603, 3b605, 3b606]: 137 mg, Selenium [3b801, 3b811, 3b812]: 0.07 mg - Preservatives - Antioxidants.

*** Values reflect only levels added to the formula, not those naturally occurring in components of the diet.



URINARY S/O + HYPOALLERGENIC



Available in 3.5, 10 and 15 kg bags

RECOMMENDED DAILY INTAKE & FEEDING DURATION

ADULT DOG	DOG'S WEIGHT (kg)	BODY CONDITION		
		THIN	NORMAL	OVERWEIGHT
		grams	grams	grams
	2	57	50	43
	4	95	84	72
	6	129	114	98
	8	160	141	122
	10	189	167	144
	15	257	226	195
	20	318	280	242
	25	376	331	286
	30	432	380	328
	35	485	426	368
	40	536	471	407
	45	585	515	445
	50	633	557	481
	55	680	598	517
	60	726	639	552
	70	815	717	619
	80	901	793	685

ROYAL CANIN® URINARY S/O + HYPOALLERGENIC dry is a complete dietetic feed for dogs formulated for the dissolution of struvite stones, the reduction of struvite stone recurrence, and for the reduction of ingredient and nutrient intolerances. Urine undersaturating or metastabilising properties for struvite and/or Urine acidifying properties. Hydrolysed protein sources and selected carbohydrate sources.

RECOMMENDATIONS: It is recommended that advice from a veterinarian be sought before use and before extending the period of use. Feed URINARY S/O + HYPOALLERGENIC dry for 5 to 12 weeks for the dissolution of struvite stones, initially up to 6 months for the reduction of struvite stone recurrence, and 3 to 8 weeks for the reduction of ingredient and nutrient intolerances: if signs of intolerance disappear this feed can be used initially up to one year.



This diet is formulated to promote a healthy urinary environment.



RECOMMENDED FOR CASES OF

Urinary disorders:

- Struvite uroliths: dissolution and management of recurrence
- Calcium oxalate uroliths: management of recurrence

Dermatological/GI conditions:

- Adverse Food Reactions (AFR)
- Canine Atopic Dermatitis (CAD)
- Chronic diarrhoea
- Inflammatory Bowel Disease (IBD)
- Exocrine Pancreatic Insufficiency (EPI)

NOT RECOMMENDED FOR CASES OF

- Chronic Kidney Disease (CKD)
- Concurrent use of urine-acidifying medications
- Growth, gestation/lactation
- Cardiac disease (when sodium restriction is required) / Heart failure



URINE DILUTION

Urine dilution makes the urine less liable to form crystals.



LOW RSS

A low concentration of ions helps the dissolution and limits the formation of common urinary stones.



HYDROLYSED PROTEIN

Hydrolysed protein with low molecular weight to ensure the food is hypoallergenic.



SKIN BARRIER

Formulated to support the skin's natural protective barrier for optimal skin health.

KEY VALUES

	Per 1000Kcal	Per 100g as fed
Moisture	25.59 g	9.5 g
Protein	53.88 g	20 g
Fat content	40.41 g	15 g
Crude Fibre	6.47 g	2.4 g
Ash	22.36 g	8.3 g
Carbohydrate (NFE)	120.69 g	44.8 g
Dietary Fibre	18.59 g	6.9 g
Omega-6	9.56 g	3.55 g
Omega-3	2.16 g	0.8 g
EPA + DHA	0.97g	0.36 g
Calcium	1.51 g	0.56 g
Phosphorus	1.35 g	0.5 g
Potassium	2.16 g	0.8 g
Sodium	3.37 g	1.25 g
Chloride	4.98 g	1.85 g
Magnesium	0.11 g	0.04 g
Copper	4.04 mg	1.5 mg
Metabolisable energy*		371.2 Kcal

*Calculated according to NRC 2006 CF.
Sulphur 0.6%.

COMPOSITION

Rice flour, hydrolysed soya protein isolate, animal fats, rice, minerals, chicory pulp, soya oil, vegetable fibres, fish oil, fructo-oligosaccharides, sodium butyrate, borage oil, mangold meal.

Protein source: hydrolysed soya protein isolate (17.5%). Carbohydrate sources: rice flour (38.9%), rice (10.1%).

ADDITIVES (KG)**

Nutritional additives: Vitamin A: 30 500 IU, Vitamin D3: 800 IU, Iron (3b103): 41 mg, Iodine (3b201, 3b202): 3.7 mg, Copper (3b405, 3b406): 15 mg, Manganese (3b502, 3b504): 54 mg, Zinc (3b603, 3b605, 3b606): 160 mg, Selenium (3b801, 3b811, 3b812): 0.26 mg - Technological additives: Clinoptilolite of sedimentary origin: 10 g - Preservatives - Antioxidants.

** Values reflect only levels added to the formula, not those naturally occurring in components of the diet.



HYPOALLERGENIC + SATIETY



Available in 3 and 10 kg bags

RECOMMENDED DAILY INTAKE & FEEDING DURATION

	IDEAL WEIGHT (kg)	STAGE			
		WEIGHT LOSS		MAINTENANCE AFTER WEIGHT LOSS	
		grams	cups	grams	cups
ADULT DOG	2	38	3/8	41	4/8
	4	63	6/8	70	6/8
	6	86	1	94	1
	8	106	1 + 2/8	117	1 + 3/8
	10	126	1 + 3/8	138	1 + 4/8
	15	170	1 + 7/8	187	2 + 1/8
	20	211	2 + 3/8	232	2 + 5/8
	25	250	2 + 7/8	275	3 + 1/8
	30	286	3 + 2/8	315	3 + 4/8
	35	322	3 + 5/8	354	4
	40	355	4	391	4 + 3/8
	45	388	4 + 3/8	427	4 + 6/8
	50	420	4 + 6/8	462	5 + 2/8
	55	451	5 + 1/8	496	5 + 5/8
	60	482	5 + 3/8	530	6
	70	541	6 + 1/8	595	6 + 6/8
	80	598	6 + 6/8	658	7 + 3/8

Weight loss: Individual target weight is determined and then feeding amount established. The recommended daily intake is based on the dog's ideal weight. The starting ration may need to be adjusted during follow-up visits, based on each individual's response and the rate of weight loss. An objective of losing 1-3% bodyweight per week is generally achievable over the first 3 months, and of 0.5-2% afterwards. **Maintenance after weight loss:** Energy requirements to maintain stable body weight after weight loss are lower than those of dogs that have never had obesity. Based on current scientific knowledge, our maintenance after weight loss ratios are calculated by increasing the weight loss ration by about 10%.

It is recommended to divide the daily intake into at least 2 meals and have fresh water available at all times.

When a food allergy or a food intolerance is suspected, the dog's diet may be changed without any food transition; otherwise a classical transition can be done over 4-5 days.

ROYAL CANIN® HYPOALLERGENIC + SATIETY dry is a complete dietetic feed for dogs formulated for the reduction of ingredient and nutrient intolerances and for the reduction of excessive body weight. Hydrolysed protein sources and selected carbohydrate sources.

RECOMMENDATIONS: your advice as a veterinarian is necessary before use and before extending the period of use of this diet. For the reduction of ingredient and nutrient intolerances, feed 3 to 8 weeks; if signs of intolerance disappear this feed can be used initially up to one year before recheck. Feed until target body weight is achieved and after if required to maintain target body weight. For an efficient weight loss or ideal weight maintenance, the recommended daily energy intake should not be exceeded. The diet may be continued for the dog's lifetime with vet follow-up.

RECOMMENDED FOR CASES OF

Dermatological/GI conditions:

- Adverse Food Reactions (AFR)
- Canine Atopic Dermatitis (CAD)
- Chronic Enteropathy, including Inflammatory Bowel Disease (IBD)
- GI conditions requiring a high fibre level
- Fibre-responsive colitis/diarrhoea (including stress diarrhoea)

+

Weight management:

- Overweight/obesity: weight loss and maintenance after weight loss
- Stabilised diabetes mellitus with concomitant overweight/obesity
- Hyperlipidaemia with concomitant overweight/obesity

NOT RECOMMENDED FOR CASES OF

- Growth, gestation or lactation
- Conditions where high fibre content is not recommended (eg: chronic constipation, severe arthritis making defecation difficult)



HYDROLYSED PROTEIN

Hydrolysed protein with low molecular weight to ensure the food is hypoallergenic.



SKIN BARRIER

Formulated to support the skin's natural protective barrier for optimal skin health.



EFFECTIVE WEIGHT MANAGEMENT

Provides safe weight loss and helps avoid weight regain. 97% of dogs lost weight in 3 months using this nutritional approach*



BEGGING CONTROL

High level of selected fibres helps to keep dogs satisfied between meals. Nutritional approach shown to help control begging in 83% of dogs during weight loss*

*Flanagan J. et al (2017) Success of a weight loss plan for overweight dogs: The results of an international weight loss study. PLoS ONE 12(9): e0184199.

KEY VALUES

	Per 1000 kcal	Per 100g as fed
Moisture	36.74 g	9.50 g
Protein	116 g	30 g
Fat content	36.74 g	9.50 g
Crude Fibre	64.19 g	16.60 g
Ash	22.04 g	5.70 g
Carbohydrate (NFE)	111 g	28.70 g
Dietary Fibre	108 g	28 g
Omega-6 fatty acids	8.43 g	2.18 g
Omega-3 fatty acids	2.90 g	0.75 g
EPA + DHA	1.59 g	0.41 g
Calcium	3.05 g	0.79 g
Phosphorus	2.32 g	0.60 g
Potassium	3.09 g	0.80 g
Sodium	1.78 g	0.46 g
Chloride	2.59 g	0.67 g
Magnesium	0.43 g	0.11 g
Copper	7.35 mg	1.90 mg
Metabolisable energy**		258.60 kcal

**Calculated according to NRC 2006 CF.

COMPOSITION

Hydrolysed soya protein isolate***, cellulose powder, rice****, hydrolysed poultry liver, minerals, animal fats, beet pulp, fish oil, soya oil, psyllium husks and seeds, sodium butyrate, fructo-oligosaccharides, mono- and diglycerides of palmitic and stearic acids esterified with citric acid, marigold meal, borage oil. ***Protein source: 30.5% ****Carbohydrate source: 22.5%.

ADDITIVES (KG)*****

Nutritional additives: Vitamin A: 3360 IU, Vitamin D3: 700 IU, Iron (3b103): 38 mg, Iodine (3b201, 3b202): 3.4 mg, Copper (3b405, 3b406): 14 mg, Manganese (3b502, 3b504): 50 mg, Zinc (3b603, 3b605, 3b606): 148 mg, Selenium (3b801, 3b811, 3b812): 0.24 mg - Preservatives - Antioxidants.

***** Values reflect only levels added to the formula, not those naturally occurring in components of the diet.





HYPOALLERGENIC + SATIETY SMALL DOGS



Available in 1 and 3 kg bags

RECOMMENDED DAILY INTAKE & FEEDING DURATION

ADULT DOG	IDEAL WEIGHT (kg)	STAGE			
		WEIGHT LOSS		MAINTENANCE AFTER WEIGHT LOSS	
		grams	cups	grams	cups
	2	38	3/8	41	4/8
	2.5	44	4/8	49	4/8
	3	51	4/8	56	5/8
	3.5	57	5/8	63	6/8
	4	63	6/8	70	6/8
	4.5	69	6/8	76	7/8
	5	75	7/8	82	7/8
	5.5	80	7/8	88	1
	6	86	1	94	1
	6.5	91	1	100	1 + 1/8
	7	96	1	106	1 + 1/8
	7.5	101	1 + 1/8	111	1 + 2/8
	8	106	1 + 1/8	117	1 + 2/8
	8.5	111	1 + 2/8	122	1 + 3/8
	9	116	1 + 2/8	128	1 + 3/8
	9.5	121	1 + 3/8	133	1 + 4/8
	10	126	1 + 3/8	138	1 + 4/8

Weight loss: Individual target weight is determined and then feeding amount established. The recommended daily intake is based on the dog's ideal weight. The starting ration may need to be adjusted during follow-up visits, based on each individual's response and the rate of weight loss. An objective of losing 1-3% bodyweight per week is generally achievable over the first 3 months, and of 0.5-2% afterwards. Maintenance after weight loss: Energy requirements to maintain stable body weight after weight loss are lower than those of dogs that have never had obesity. Based on current scientific knowledge, our maintenance after weight loss ratios are calculated by increasing the weight loss ratio by about 10%.

It is recommended to divide the daily intake into at least 2 meals and have fresh water available at all times. When a food allergy or a food intolerance is suspected, the dog's diet may be changed without any food transition; otherwise a classical transition can be done (over 4-5 days).

ROYAL CANIN® HYPOALLERGENIC - SATIETY SMALL DOGS dry is a complete dietetic feed for dogs formulated for the reduction of ingredient and nutrient intolerances and for the reduction of excessive body weight. Hydrolysed protein sources and selected carbohydrate sources.

RECOMMENDATIONS: your advice as a veterinarian is necessary before use and before extending the period of use of this diet. For the reduction of ingredient and nutrient intolerances, feed 3 to 8 weeks: if signs of intolerance disappear this feed can be used initially up to one year before recheck. Feed until target body weight is achieved and after if required to maintain target body weight. For an efficient weight loss or ideal weight maintenance, the recommended daily energy intake should not be exceeded. The diet may be continued for the dog's lifetime with vet follow-up.

RECOMMENDED FOR CASES OF

Dermatological/GI conditions:

- Adverse Food Reactions (AFR)
- Canine Atopic Dermatitis (CAD)
- Chronic Enteropathy, including Inflammatory Bowel Disease (IBD)
- GI conditions requiring a high fibre level
- Fibre responsive colitis/diarrhoea (including stress diarrhoea)

+ Weight management:

- Overweight/obesity: weight loss and maintenance after weight loss
- Stabilised diabetes mellitus with concomitant overweight/obesity
- Hyperlipidaemia with concomitant overweight/obesity

NOT RECOMMENDED FOR CASES OF

- Growth, gestation or lactation
- Conditions where high fibre content is not recommended (eg: chronic constipation, severe arthritis making defecation difficult)



HYDROLYSED PROTEIN

Hydrolysed protein with low molecular weight to ensure the food is hypoallergenic.



SKIN BARRIER

Formulated to support the skin's natural protective barrier for optimal skin health.



EFFECTIVE WEIGHT MANAGEMENT

Provides safe weight loss and helps avoid weight regain. 97% of dogs lost weight in 3 months using this nutritional approach*



BEGGING CONTROL

High level of selected fibres helps to keep dogs satisfied between meals. Nutritional approach shown to help control begging in 83% of dogs during weight loss*



SMALL KIBBLE

Carefully calibrated kibble size to fit small jaws.

*Flanagan J. et al (2017) Success of a weight loss plan for overweight dogs: The results of an international weight loss study. PLoS ONE 12(9): e0184199.

KEY VALUES

	Per 1000 kcal	Per 100g as fed
Moisture	36.74 g	9.50 g
Protein	116 g	30 g
Fat content	36.74 g	9.50 g
Crude Fibre	64.19 g	16.60 g
Ash	22.04 g	5.70 g
Carbohydrate (NFE)	111 g	28.70 g
Dietary Fibre	108 g	28 g
Omega-6 fatty acids	8.43 g	2.18 g
Omega-3 fatty acids	2.90 g	0.75 g
EPA + DHA	1.59 g	0.41 g
Calcium	3.05 g	0.79 g
Phosphorus	2.32 g	0.60 g
Potassium	3.09 g	0.80 g
Sodium	1.78 g	0.46 g
Chloride	2.59 g	0.67 g
Magnesium	0.43 g	0.11 g
Copper	7.35 mg	1.90 mg
Metabolisable energy**		258.60 kcal

**Calculated according to NRC 2006 CF.

Essential Fatty Acid (Linoleic Acid): 2.08%

COMPOSITION

Hydrolysed soya protein isolate***, cellulose powder, rice****, hydrolysed poultry liver, minerals, animal fats, beet pulp, fish oil, soya oil, psyllium husks and seeds, sodium butyrate, fructo-oligosaccharides, mono- and diglycerides of palmitic and stearic acids esterified with citric acid, marigold meal, borage oil.

Protein source: 30.4%, *Carbohydrate source: 22.6%.

ADDITIVES (KG)*****

Nutritional additives: Vitamin A: 33300 IU, Vitamin D3: 700 IU, Iron (3b103): 38 mg, Iodine (3b201, 3b202): 3.4 mg, Copper (3b405, 3b406): 14 mg, Manganese (3b502, 3b504): 50 mg, Zinc (3b603, 3b605, 3b606): 146 mg, Selenium (3b801, 3b811, 3b812): 0.24 mg - Preservatives - Antioxidants.

***** Values reflect only levels added to the formula, not those naturally occurring in components of the diet.





HEPATIC

Gastrointestinal



Available in 1.5 kg, 6kg

RECOMMENDED DAILY INTAKE

	DOG'S WEIGHT (kg)	WEIGHT CONDITION		
		UNDERWEIGHT	IDEAL WEIGHT	OVERWEIGHT
		grams	grams	grams
ADULT DOG	2	54	48	41
	4	91	80	69
	6	123	109	94
	8	153	135	116
	10	181	159	138
	15	245	216	187
	20	305	268	232
	25	360	317	274
	30	413	363	314
	35	463	408	352
	40	512	451	389
	45	560	492	425
	50	606	533	460

It is recommended to divide the daily intake into several small meals throughout the day and have fresh water available at all times.

For acute cases, feeding a GI diet at least 2-3 weeks (continuing after clinical signs have resolved) helps support optimal pet recovery and regeneration of the intestinal villi. If signs have not resolved in 5-7 days, consider additional diagnostics such as blood tests, imaging and infectious disease testing. For liver diseases, feeding duration varies depending on the aetiology and on the regenerative ability of the hepatic tissue.

For porto-systemic shunts and hepatic encephalopathies, the nutritional goal is to feed the pet the highest level of protein that they can tolerate, preferably from vegetable or dairy sources. If signs of encephalopathy persist, further protein restriction is warranted and the dog should be switched to ROYAL CANIN® HEPATIC diets.

ROYAL CANIN® HEPATIC GASTROINTESTINAL dry is a complete dietetic feed for dogs formulated for the compensation for malabsorption, for the reduction of intestinal absorptive disorders, for the support of liver function in the case of chronic liver insufficiency and for the reduction of copper in the liver. Highly digestible feed with increased sodium and potassium. Moderate level of protein, selected and highly digestible. Restricted level of copper.

RECOMMENDATIONS: your advice as a veterinarian is necessary before use and before extending the period of use. For the compensation for malabsorption and for the reduction of intestinal absorptive disorders, feed this diet initially up to 12 weeks before recheck and lifetime in the case of chronic pancreatic insufficiency. Feed this diet initially up to 4 months for the support of liver function in the case of chronic liver insufficiency and initially up to 6 months for the reduction of copper in the liver before recheck. The diet may be continued for the dog's lifetime with vet follow-up.

PROVIDES DIGESTIVE AND LIVER SUPPORT.

RECOMMENDED FOR CASES OF

- Copper storage disease*
- Portosystemic shunt*
- Acute hepatopathy
- Chronic hepatitis*
- Hepatic encephalopathy*
- Acute and chronic diarrhoea
- Maldigestion/malabsorption
- Chronic Inflammatory Enteropathy (CIE) including Inflammatory Bowel Disease (IBD)

NOT RECOMMENDED IN CASE OF

- Gestation or lactation
- Pancreatitis or history of pancreatitis
- Hyperlipidemia
- Lymphangiectasia

*In case of severe hepatic encephalopathy, a more protein-restricted diet might be needed (Hepatic)

For puppies with shunts, can be used under vet supervision, with close follow-up and monitoring of growth.



ADAPTED PROTEIN CONTENT

Moderate levels of high-quality proteins to support liver function.



LOW COPPER

Low level of copper to help minimise its accumulation in liver cells.



DIGESTIVE SUPPORT

A highly digestible formula with balanced fibres, including prebiotics, to support a healthy digestion and microbiota.



HIGH ENERGY

High energy content to reduce meal volume and help compensate for a decreased appetite.

KEY VALUES

	Per 1000 kcal	Per 100 g as fed
Moisture	24.48 g	9.50 g
Protein	51.53 g	20 g
Fat content	41.23 g	16 g
Crude Fibre	5.15 g	2 g
Ash	14.69 g	5.70 g
Carbohydrate (NFE)	121 g	46.80 g
Total Dietary Fiber (TDF)	19.07 g	7.40 g
Omega-6 fatty acids	9.15 g	3.55 g
Omega-3 fatty acids	2.11 g	0.82 g
EPA + DHA	1.03 g	0.40 g
Calcium	1.88 g	0.73 g
Phosphorus	1.42 g	0.55 g
Potassium	2.11 g	0.82 g
Sodium	0.82 g	0.32 g
Chloride	2.37 g	0.92 g
Magnesium	0.23 g	0.09 g
Copper	0.80 mg	0.31 mg
Zinc	43.80 mg	17 mg
Metabolisable energy*		388.1

*Calculated according to NRC 2006 CF

COMPOSITION

Broken rice**, animal fats**, hydrolysed soya protein isolate*, maize**, wheat gluten*, beet pulp, animal protein hydrolysate**, minerals, refined soya oil**, vegetable fibres, fish oil**, psyllium husks, fructo-oligosaccharides, algal oil Schizochytrium sp. (source of EPA+DHA)**, yeast components (source of manno-oligosaccharides 0.05%), marigold meal.

*Protein sources: 16.9%. **Highly digestible ingredients: 86.6%.

ADDITIVES (KG)**

Nutritional additives: Vitamin A: 19700 IU, Vitamin D3: 1200 IU, Vitamin E: 677 mg, Vitamin C: 349 mg, Iron (3b103): 118 mg, Iodine (3b201, 3b202): 5.7 mg, Manganese (3b502, 3b505): 55 mg, Zinc (3b603, 3b605, 3b612): 149 mg, Selenium (3b801, 3b811, 3b812): 0.39 mg - Technological additives: Clnoptilolite of sedimentary origin: 5 g - Preservatives - Antioxidants.

**Values reflect only levels added to the formula, not those naturally occurring in components of the diet.



VET SERVICES

Scan to access the Vet Services Feeding Amount Calculator for a precise, individualized feeding recommendation.



HEPATIC

Gastrointestinal

PROVIDES DIGESTIVE AND LIVER SUPPORT.

RECOMMENDED FOR CASES OF

- Copper storage disease*
- Portosystemic shunt*
- Acute hepatopathy
- Chronic hepatitis*
- Hepatic encephalopathy*
- Acute and chronic diarrhoea
- Maldigestion/malabsorption
- Chronic Inflammatory Enteropathy (CIE) including Inflammatory Bowel Disease (IBD)

NOT RECOMMENDED IN CASE OF

- Gestation or lactation
- Pancreatitis or history of pancreatitis
- Hyperlipidemia
- Lymphangiectasia

*In case of severe hepatic encephalopathy, a more protein-restricted diet might be needed (Hepatic)



Available in 200g

RECOMMENDED DAILY INTAKE

ADULT DOG	DOG'S WEIGHT (kg)	WEIGHT CONDITION					
		UNDERWEIGHT		IDEAL WEIGHT		OVERWEIGHT	
		grams	can	grams	can	grams	can
	2	143	3/4	125	3/4	108	2/4
	4	240	1+1/4	211	1	182	1
	6	325	1+3/4	286	1+2/4	247	1+1/4
	8	403	2	355	1+3/4	306	1+2/4
	10	477	2+2/4	419	2	362	1+3/4
	15	646	3+1/4	568	2+3/4	491	2+2/4
	20	801	4	705	3+2/4	609	3
	25	947	4+3/4	834	4+1/4	720	3+2/4
	30	1086	5+2/4	956	4+3/4	826	4+1/4
	35	1219	6	1073	5+1/4	927	4+3/4
	40	1348	6+3/4	1186	6	1024	5
	45	1472	7+1/4	1296	6+2/4	1119	5+2/4
	50	1593	8	1402	7	1211	6

It is recommended to divide the daily intake into several small meals throughout the day and have fresh water available at all times.

For acute cases, feeding a GI diet at least 2-3 weeks (continuing after clinical signs have resolved) helps support optimal pet recovery and regeneration of the intestinal villi. If signs have not resolved in 5-7 days, consider additional diagnostics such as blood tests, imaging and infectious disease testing. For liver diseases, feeding duration varies depending on the aetiology and on the regenerative ability of the hepatic tissue.

For porto-systemic shunts and hepatic encephalopathies, the nutritional goal is to feed the pet the highest level of protein that they can tolerate, preferably from vegetable or dairy sources. If signs of encephalopathy persist, further protein restriction is warranted and the dog should be switched to ROYAL CANIN® HEPATIC diets.

ROYAL CANIN HEPATIC GASTROINTESTINAL loaf is a complete dietetic feed for dogs formulated for the support of liver function in the case of chronic liver insufficiency and for the reduction of copper in the liver. Moderate level of protein, selected and highly digestible. Restricted level of copper.

RECOMMENDATIONS: your advice as a veterinarian is necessary before use and before extending the period of use. Feed this diet initially up to 4 months for the support of liver function in the case of chronic liver insufficiency and initially up to 6 months for the reduction of copper in the liver before recheck. The diet may be continued for the dog's lifetime with vet follow-up.



ADAPTED PROTEIN CONTENT

Moderate levels of high-quality proteins to support liver function.



LOW COPPER

Low level of copper to help minimise its accumulation in liver cells.



DIGESTIVE SUPPORT

A highly digestible formula with balanced fibres, including prebiotics, to support a healthy digestion and microbiota.



HIGH ENERGY

High energy content to reduce meal volume and help compensate for a decreased appetite.

KEY VALUES

	Per 1000 kcal	Per 100 g as fed
Moisture	450 g	64.50 g
Protein	51.64 g	7.40 g
Fat content	27.91 g	4 g
Crude Fibre	5.58 g	0.80 g
Ash	11.86 g	1.7 g
Carbohydrate (NFE)	151 g	21.60 g
Total Dietary Fiber (TDF)	16.05 g	2.30 g
Omega-6 fatty acids	7.68 g	1.10 g
Omega-3 fatty acids	1.61 g	0.23 g
EPA + DHA	1.12 g	0.16 g
Calcium	2.09 g	0.30 g
Phosphorus	1.61 g	0.23 g
Potassium	1.47 g	0.21 g
Sodium	0.49 g	0.07 g
Chloride	1.05 g	0.15 g
Magnesium	0.22 g	0.03 g
Copper	1.19 mg	0.17 mg
Zinc	48.85 mg	7 mg
Metabolisable energy*		143.3 kcal

*Calculated according to NRC 2006 CF

COMPOSITION Off-pack information can differ from on-pack

Meat and animal derivatives*, cereals**, vegetable protein extracts*, derivatives of vegetable origin, oils and fats**, minerals, yeasts, various sugars**.

*Protein sources: 29.3%. **Highly digestible ingredients: 56.4%.

ADDITIVES (KG)***

Nutritional additives: Vitamin D3: 2770 IU, Vitamin E: 272 mg, Vitamin C: 219 mg, Iron (3b103): 23 mg, Iodine (3b202): 0.68 mg, Copper (3b405, 3b407): 0.5 mg, Manganese (3b502, 3b503, 3b505): 9.1 mg, Zinc (3b603, 3b612): 57 mg, Selenium (3b801, 3b811, 3b812): 0.04 mg - Technological additives: Clinoptilolite of sedimentary origin: 0.2 g.

***Values reflect only levels added to the formula, not those naturally occurring in components of the diet.



VET SERVICES

Scan to access the Vet Services Feeding Amount Calculator for a precise, individualized feeding recommendation.



GASTROINTESTINAL

Hydrolysed protein



Available in 3.5 kg, 9 kg

RECOMMENDED DAILY INTAKE

ADULT DOG	DOG'S WEIGHT (kg)	WEIGHT CONDITION		
		UNDERWEIGHT	IDEAL WEIGHT	OVERWEIGHT
		grams	grams	grams
	2	54	47	41
	4	90	80	69
	6	122	108	93
	8	152	134	115
	10	180	158	137
	15	243	214	185
	20	302	266	230
	25	357	314	271
	30	409	360	311
	35	460	405	349
	40	508	447	386
	45	555	488	422
	50	601	529	457

It is recommended to divide the daily intake into several small meals throughout the day and have fresh water available at all times.

For acute cases, feeding a GI diet at least 2-3 weeks (continuing after clinical signs have resolved) helps support optimal pet recovery and regeneration of the intestinal villi. If signs have not resolved in 5-7 days, consider additional diagnostics such as blood tests, imaging and infectious disease testing.

For chronic enteropathies, nutritional management is key. Successive 2-week dietary trials after a 7-10 day transition should be conducted until the most effective diet is identified. After a 2-week trial is completed: if full remission achieved, continue diet long term, in case of partial remission, continue trial for 2 more weeks and if there is no improvement, try a different GI diet. If no diet leads to clinical remission, the most effective diet at reducing signs should be continued long term alongside appropriate medications.

ROYAL CANIN® GASTROINTESTINAL HYDROLYSED PROTEIN dry is a complete dietetic feed for dogs formulated for the reduction of ingredient and nutrient intolerances, for the compensation for maldigestion and for the reduction of intestinal absorptive disorders. Hydrolysed protein sources and selected carbohydrate sources. Highly digestible feed with increased sodium and potassium.

RECOMMENDATIONS: your advice as a veterinarian is necessary before use and before extending the period of use of this diet. For the reduction of ingredient and nutrient intolerances, feed this diet 3 to 8 weeks before recheck and up to one year before reassessing nutritional recommendation if signs of intolerance disappear. For the compensation for maldigestion and for the reduction of intestinal absorptive disorders, feed this diet initially up to 12 weeks before recheck and lifetime in the case of chronic pancreatic insufficiency. The diet may be continued for the dog's lifetime with vet follow-up.

PROVIDES DIGESTIVE SUPPORT WHILE MINIMISING THE RISK OF ADVERSE FOOD REACTIONS.

RECOMMENDED FOR CASES OF

- Chronic Inflammatory Enteropathy (CIE) - formerly called Inflammatory Bowel Disease (IBD)
- Acute and chronic diarrhoea
- Maldigestion/malabsorption including Exocrine Pancreatic Insufficiency (EPI)
- Adverse Food Reactions (AFR)
- Impaired appetite, feeding post surgery
- Acute hepatopathy without hepatic encephalopathy

NOT RECOMMENDED IN CASE OF

- Lymphangiectasia¹
- Hyperlipidemia¹
- Pancreatitis or history of pancreatitis¹
- Growth, gestation or lactation²
- Hepatic encephalopathy²

¹ In these cases the recommended diet is Gastrointestinal Low Fat Hypoallergenic (or other Gastrointestinal Low Fat diets) - ² For puppies or gestating/lactating dams, the most appropriate is Gastrointestinal Puppy ³ In these cases the recommended diet are Hepatic Gastrointestinal or Hepatic

For puppies, can be used under vet supervision, with close follow-up and monitoring of growth.



DIGESTIVE SUPPORT

A highly digestible formula with balanced fibres to support a healthy digestion and transit.



MICROBIOME SUPPORT

Formula with selected prebiotics to support a healthy gut and intestinal microbiome.



HIGH ENERGY

High energy content to reduce meal volume and help compensate for a decreased appetite.



HYDROLYSED PROTEIN

Highly digestible hydrolysed protein to ensure the food is hypoallergenic.

KEY VALUES

	Per 1000 kcal	Per 100 g as fed
Moisture	23 g	9 g
Protein	63.89 g	25 g
Fat content	51.11 g	20 g
Crude Fibre	10.73 g	4.20 g
Ash	16.10 g	6.3 g
Carbohydrate (NFE)	90.72 g	35.50 g
Total Dietary Fiber (TDF)	27.09 g	10.60 g
Omega-6 fatty acids	10.58 g	4.14 g
Omega-3 fatty acids	2.43 g	0.95 g
EPA + DHA	1.02 g	0.40 g
Calcium	2.04 g	0.80 g
Phosphorus	1.53 g	0.60 g
Potassium	1.74 g	0.68 g
Sodium	1.05 g	0.41 g
Chloride	1.97 g	0.77 g
Magnesium	0.23 g	0.09 g
Copper	4.88 mg	1.91 mg
Zinc	43.44 mg	17 mg
Metabolisable energy*		391.3 kcal

*Calculated according to NRC 2006 CF - Essential fatty acid (linoleic acid): 4.02%

COMPOSITION

Hydrolysed soya protein isolate*, broken rice**, animal fats***, rice flour**, animal protein hydrolysate***, vegetable fibres, beet pulp, minerals, fish oil***, refined soy oil***, fructo-oligosaccharides, psyllium husks, yeast components (source of manno-oligosaccharides 0.05%), marigold meal.

*Protein source: 23%. **Carbohydrate sources: 36.3%. ***Highly digestible ingredients: 82.1%.

ADDITIVES (KG)**

Vitamin A: 34800 IU, Vitamin D3: 800 IU, Vitamin E: 820 mg, Vitamin C: 374 mg, Iron (3b103): 39 mg, Iodine (3b201, 3b202): 3.6 mg, Copper (3b405): 14.6 mg, Manganese (3b502, 3b505): 52 mg, Zinc (3b603, 3b605, 3b612): 153 mg, Selenium (3b801, 3b811, 3b812): 0.25 mg - Technological additives: Cinoptilolite of sedimentary origin: 10 g - Preservatives - Antioxidants.

**Values reflect only levels added to the formula, not those naturally occurring in components of the diet.



VET SERVICES

Scan to access the Vet Services Feeding Amount Calculator for a precise, individualized feeding recommendation.



GASTROINTESTINAL

Hydrolysed protein



This diet supports a healthy urinary environment.

S/O INDEX

Available in 400g

RECOMMENDED DAILY INTAKE

	DOG'S WEIGHT (kg)	WEIGHT CONDITION					
		UNDERWEIGHT		IDEAL WEIGHT		OVERWEIGHT	
		grams	can	grams	can	grams	can
ADULT DOG	2	211	2/4	185	2/4	160	2/4
	4	354	1	312	3/4	269	3/4
	6	480	1+1/4	423	1	365	1
	8	596	1+2/4	524	1+1/4	453	1+1/4
	10	704	1+3/4	620	1+2/4	535	1+1/4
	15	955	2+2/4	840	2	726	1+3/4
	20	1185	3	1042	2+2/4	900	2+1/4
	25	1400	3+2/4	1232	3	1064	2+3/4
	30	1606	4	1413	3+2/4	1220	3
	35	1802	4+2/4	1586	4	1370	3+2/4
	40	1992	5	1753	4+2/4	1514	3+3/4
	45	2176	5+2/4	1915	4+3/4	1654	4+1/4
	50	2355	6	2072	5+1/4	1790	4+2/4

It is recommended to divide the daily intake into several small meals throughout the day and have fresh water available at all times.

For acute cases, feeding a GI diet at least 2-3 weeks (continuing after clinical signs have resolved) helps support optimal pet recovery and regeneration of the intestinal villi. If signs have not resolved in 5-7 days, consider additional diagnostics such as blood tests, imaging and infectious disease testing.

For chronic enteropathies, nutritional management is key. Successive 2-week dietary trials after a 7-10 day transition should be conducted until the most effective diet is identified. After a 2-week trial is completed: if full remission achieved, continue diet long term; in case of partial remission, continue trial for 2 more weeks and if there is no improvement, try a different GI diet. If no diet leads to clinical remission, the most effective diet at reducing signs should be continued long term alongside appropriate medications.

ROYAL CANIN® GASTROINTESTINAL HYDROLYSED PROTEIN (Loaf) is a complete dietetic feed for dogs formulated for the reduction of ingredient and nutrient intolerances, for the compensation for maldigestion and for the reduction of intestinal absorptive disorders. Hydrolysed protein sources and selected carbohydrate sources. Highly digestible feed with increased sodium and potassium.

RECOMMENDATIONS: your advice as a veterinarian is necessary before use and before extending the period of use of this diet. For the reduction of ingredient and nutrient intolerances, feed this diet 3 to 8 weeks before recheck and up to one year before reassessing nutritional recommendation if signs of intolerance disappear. For the compensation for maldigestion and for the reduction of intestinal absorptive disorders, feed this diet initially up to 12 weeks before recheck and lifetime in the case of chronic pancreatic insufficiency. The diet may be continued for the dog's lifetime with vet follow-up.

PROVIDES DIGESTIVE SUPPORT WHILE MINIMISING THE RISK OF ADVERSE FOOD REACTIONS.

RECOMMENDED FOR CASES OF

- Chronic Inflammatory Enteropathy (CIE) - formerly Inflammatory Bowel Disease (IBD)
- Acute and chronic diarrhoea
- Maldigestion/malabsorption including Exocrine Pancreatic Insufficiency (EPI)
- Adverse Food Reactions (AFR)
- Impaired appetite, feeding post surgery
- Acute hepatopathy without hepatic encephalopathy

NOT RECOMMENDED IN CASE OF

- Lymphangiectasia¹
- Hyperlipidemia¹
- Pancreatitis or history of pancreatitis¹
- Growth, gestation or lactation²
- Hepatic encephalopathy²

1. In these cases the recommended diet is Gastrointestinal Low Fat Hypoallergenic (or other Gastrointestinal Low Fat diets) - 2. For puppies or gestating/lactating dams, the most appropriate is Gastrointestinal Puppy 3. In these cases the recommended diet are Hepatic Gastrointestinal or Hepatic.

For puppies, can be used under vet supervision, with close follow-up and monitoring of growth.



DIGESTIVE SUPPORT

A highly digestible formula with balanced fibres to support a healthy digestion and transit.



MICROBIOME SUPPORT

Formula with selected prebiotics to support a healthy gut and intestinal microbiome.



HIGH ENERGY

High energy content to reduce meal volume and help compensate for a decreased appetite.



HYDROLYSED PROTEIN

Highly hydrolysed protein to ensure the food is hypoallergenic.

KEY VALUES

	Per 1000 kcal	Per 100 g as fed
Moisture	798g	73.70 g
Protein	69.34g	6.40 g
Fat content	50.92 g	4.70 g
Crude Fibre	33.59 g	3.10 g
Ash	26 g	2.40 g
Carbohydrate (NFE)	105 g	9.70 g
Total Dietary Fiber (TDF)	49.84 g	4.60 g
Omega-6 fatty acids	24.92 g	2.30 g
Omega-3 fatty acids	1.73 g	0.16 g
EPA + DHA	1.19 g	0.11 g
Calcium	3.68 g	0.34 g
Phosphorus	2.82 g	0.26 g
Potassium	2.17 g	0.20 g
Sodium	2.28 g	0.21 g
Chloride	2.06 g	0.19 g
Magnesium	0.23 g	0.02 g
Copper	4.88 mg	0.45 mg
Zinc	36.84 mg	3.40 mg
Metabolisable energy*		92.3 kcal

*Calculated according to NRC 2006 CF - Essential fatty acid (linoleic acid): 2.20%.

COMPOSITION Off-pack information can differ from on-pack

Pea starch**, hydrolysed soya protein isolate*, powder cellulose, refined sunflower oil***, animal protein hydrolysate*, minerals, fish oil***, yeast components, marigold meal.

*Protein sources: 6.7% - **Carbohydrate source: 8.5% - ***Highly digestible ingredients: 19.3%

ADDITIVES (KG)***

Vitamin A: 6000 IU, Vitamin D3: 270 IU, Vitamin E: 202 mg, Vitamin C: 104 mg, Taurine: 1.6 g, Iron (3b103): 10 mg, Iodine (3b202): 0.87 mg, Copper (3b405, 3b407): 3.6 mg, Manganese (3b502, 3b503, 3b505): 12.8 mg, Zinc (3b603, 3b612): 29 mg, Selenium (3b801, 3b811, 3b812): 0.07 mg - Technological additives: Clnophtolite of sedimentary origin: 2 g - Sensory additives: Flavours.

***Values reflect only levels added to the formula, not those naturally occurring in components of the diet.



VET SERVICES

Scan to access the Vet Services Feeding Amount Calculator for a precise, individualized feeding recommendation.



GASTROINTESTINAL

Low fat

Hypoallergenic



Available in 2.5kg, 10kg

RECOMMENDED DAILY INTAKE

ADULT DOG	DOG'S WEIGHT (kg)	WEIGHT CONDITION		
		UNDERWEIGHT	IDEAL WEIGHT	OVERWEIGHT
		grams	grams	grams
	2	60	53	46
	4	101	89	77
	6	137	120	104
	8	169	149	129
	10	200	176	152
	15	272	239	206
	20	337	297	256
	25	398	351	303
	30	457	402	347
	35	513	451	390
	40	567	499	431
	45	619	545	471
	50	670	590	509

It is recommended to divide the daily intake into several small meals throughout the day and have fresh water available at all times. When a food allergy or a food intolerance is suspected, the dog's diet may be changed without any food transition; otherwise a classical transition can be done (over 4-5 days).

For acute cases, feeding a GI diet at least 2-3 weeks (continuing after clinical signs have resolved) helps support optimal pet recovery and regeneration of the intestinal villi. If signs have not resolved in 5-7 days, consider additional diagnostics such as blood tests, imaging and infectious disease testing.

For chronic enteropathies, nutritional management is key. Successive 2-week dietary trials after a 7-10 day transition should be conducted until the most effective diet is identified. After a 2-week trial is completed: if full remission achieved, continue diet long term, in case of partial remission, continue trial for 2 more weeks and if there is no improvement, try a different GI diet. If no diet leads to clinical remission, the most effective diet at reducing signs should be continued long term alongside appropriate medications.

ROYAL CANIN® GASTROINTESTINAL LOW FAT HYPOALLERGENIC dry is a complete dietetic feed for dogs formulated for the support of lipid metabolism in the case of hyperlipidaemia and for the reduction of ingredient and nutrient intolerances. Low level of fat. Hydrolysed protein sources and selected carbohydrate sources.

RECOMMENDATIONS: your advice as a veterinarian is necessary before use and before extending the period of use of this diet. For the reduction of ingredient and nutrient intolerances, feed this diet 3 to 8 weeks before recheck and up to one year before reassessing nutritional recommendation if signs of intolerance disappear. For the support of lipid metabolism in the case of hyperlipidaemia, feed this diet initially up to 2 months before recheck. The diet may be continued for the dog's lifetime with vet follow-up.

HIGHLY DIGESTIBLE LOW-FAT DIET FOR DOGS WITH FAT SENSITIVITY AND ADVERSE FOOD REACTIONS.

RECOMMENDED FOR CASES OF

- Chronic Inflammatory Enteropathy (CIE) - formerly called Inflammatory Bowel Disease (IBD)
- Protein-losing enteropathy (PLE)
- Acute and chronic diarrhoea
- Maldigestion/ malabsorption including Exocrine Pancreatic Insufficiency (EPI)
- Adverse Food Reactions (AFR)
- Lymphangiectasia
- Acute and chronic pancreatitis
- Hyperlipidemia

NOT RECOMMENDED IN CASE OF

- Growth, gestation or lactation



DIGESTIVE SUPPORT

A highly digestible formula with balanced fibres, including prebiotics, to support a healthy digestion and transit.



ADEQUATE ENERGY

Limited fibre content to allow provision of maintenance energy levels despite fat restriction.



LOW FAT

For the nutritional management of dogs needing a fat-restricted diet.



HYDROLYSED PROTEIN

Highly digestible hydrolysed protein to ensure the food is hypoallergenic.

KEY VALUES

	Per 1000 kcal	Per 100 g as fed
Moisture	27.08 g	9.5 g
Protein	71.27 g	25 g
Fat content	19.95 g	7 g
Crude Fibre	3.71 g	1.3 g
Ash	17.67 g	6.2 g
Carbohydrate (NFE)	145.00 g	51 g
Total Dietary Fiber (TDF)	16.53 g	5.8 g
Omega-6 fatty acids	4.59 g	1.61 g
Omega-3 fatty acids	0.91 g	0.32 g
EPA + DHA	0.43 g	0.15 g
Calcium	2.28 g	0.8 g
Phosphorus	1.71 g	0.6 g
Potassium	1.71 g	0.6 g
Sodium	1.14 g	0.4 g
Chloride	2 g	0.7 g
Magnesium	0.23 g	0.08 g
Copper	5.42 mg	1.9 mg
Zinc	48.46 mg	17 mg
Metabolisable energy*		350.8 Kcal

*Calculated according to NRC 2006 CF.

COMPOSITION

Rice flour**, hydrolysed soya protein isolate*, broken rice**, beet pulp, animal protein hydrolysate, animal fats, minerals, psyllium husks, fish oil, refined soya oil, sodium butyrate, fructo-oligosaccharides, marigold meal.

*Protein source: 21.9%. **Carbohydrate sources: 57.3%.

ADDITIVES (KG)**

Nutritional additives: Vitamin A: 34 000 IU, Vitamin D3: 800 IU, Iron (3b103): 39 mg, Iodine (3b201, 3b202): 3.5 mg, Copper (3b405): 14 mg, Manganese (3b502, 3b505): 51 mg, Zinc (3b603, 3b605, 3b612): 152 mg, Selenium (3b801, 3b811, 3b812): 0.24 mg - Technological additives: Clinoptilolite of sedimentary origin: 10 g - Preservatives - Antioxidants.

**Values reflect only levels added to the formula, not those naturally occurring in components of the diet.



VET SERVICES

Scan to access the Vet Services Feeding Amount Calculator for a precise, individualized feeding recommendation.



RENAL + HYPOALLERGENIC



Available in 4 and 10 kg bags

RECOMMENDED DAILY INTAKE & FEEDING DURATION

ADULT DOG	DOG'S WEIGHT (kg)	BODY CONDITION		
		THIN	NORMAL	OVERWEIGHT
		grams	grams	grams
	2	52 g	46 g	40 g
	4	88 g	77 g	67 g
	6	119 g	105 g	90 g
	8	148 g	130 g	112 g
	10	174 g	153 g	133 g
	15	236 g	208 g	180 g
	20	293 g	258 g	223 g
	25	347 g	305 g	263 g
	30	398 g	350 g	302 g
	35	446 g	393 g	339 g
	40	493 g	434 g	375 g
	45	539 g	474 g	409 g
	50	583 g	513 g	443 g
	55	626 g	551 g	476 g
	60	669 g	588 g	508 g
	70	750 g	660 g	570 g
	80	829 g	730 g	630 g

ROYAL CANIN® RENAL + HYPOALLERGENIC dry is a complete dietetic feed for dogs formulated for the support of renal function in the case of chronic renal insufficiency and for the reduction of ingredient and nutrient intolerance. Restricted level of phosphorus. Hydrolysed protein sources and selected carbohydrate sources.

Renal diets are recommended as soon as creatinine goes out of reference range (from IRS stage 2), before the disease further progresses, and before inappetence develops. When a food allergy or a food intolerance is suspected, the dog's diet should be changed using classical food transition. RENAL + HYPOALLERGENIC diet may be continued for the dog's lifetime.

It is recommended that a veterinarian's opinion be sought before use and before extending the period of use.

It is recommended to divide the daily intake into 2 meals.

Fresh water should be available at all times



RECOMMENDED FOR CASES OF

Renal disease

- Azotaemic Chronic Kidney Disease (IRIS stages 2 to 4)
- Chronic Kidney Disease with proteinuria (IRIS stages 1 to 4)

Dermatological/GI conditions:

- Adverse Food Reactions (AFR)
- Canine Atopic dermatitis
- Chronic diarrhoea
- Inflammatory Bowel Disease (IBD)
- Exocrine Pancreatic Insufficiency (EPI)

Other:

- Management of recurrence of urolithiasis requiring urine alkalinisation (urate and cystine) in dogs with impaired renal function and/or AFR

NOT RECOMMENDED FOR CASES OF

- Pancreatitis
- Growth, gestation/lactation



RENAL SUPPORT

Formulated with low phosphorus and moderate protein of high quality to support renal function.



ADAPTED ENERGY

Adapted energy content to reduce meal volume and help compensate for a decreased appetite.



HYDROLYSED PROTEIN

Hydrolysed protein with low molecular weight to ensure the food is hypoallergenic.



SKIN BARRIER

Formulated to support the skin's natural protective barrier for optimal skin health.

KEY VALUES

	Per 1000Kcal	Per 100g as fed
Moisture	23.57 g	9.5 g
Protein	34.73 g	14 g
Fat content	47.13 g	19 g
Crude Fibre	5.71 g	2.3 g
Ash	10.42 g	4.2 g
Carbohydrate (NFE)	126.52 g	51 g
Dietary Fibre	16.62 g	6.7 g
Omega-6	10.49 g	4.23 g
Omega-3	2.6 g	1.05 g
EPA + DHA	1.24 g	0.5 g
Calcium	0.74 g	0.3 g
Phosphorus	0.45 g	0.18 g
Sodium	0.87 g	0.35 g
Chloride	2.21 g	0.89 g
Potassium	1.49 g	0.6 g
Magnesium	0.15 g	0.06 g
Zinc	40.44 mg	16.3 mg
Vitamin D	198.46	80 IU
Vitamin A	8434.63IU	3400 IU

Metabolisable energy*

403.1 Kcal

*Calculated according to NRC 2006 CF.

Essential Fatty Acid (Linoleic Acid): 4.02 %.

COMPOSITION


Rice flour, animal fats, rice, hydrolysed soya protein isolate, chicory pulp, soya oil, fish oil, vegetable fibres, minerals, fructo-oligosaccharides, sodium butyrate, borage oil, marigold meal.

Carbohydrate sources: rice flour (44.5%), rice (12.6%). Protein source: hydrolysed soya protein isolate (8.8%).

ADDITIVES (KG)**

Nutritional additives: Vitamin A: 28500 IU, Vitamin D3: 800 IU, Iron (3b103): 39 mg, Iodine (3b201, 3b202): 3.5 mg, Copper (3b405, 3b406): 14 mg, Manganese (3b502, 3b504): 51 mg, Zinc (3b603, 3b605, 3b606): 151 mg, Selenium (3b801, 3b811, 3b812): 0.24 mg - Technological additives: Clinoptilolite of sedimentary origin: 10 g - Preservatives - Antioxidants.

** Values reflect only levels added to the formula, not those naturally occurring in components of the diet.



SATIETY + MOBILITY



Available in 1.5 kg, 3.5 kg, 9 kg bags

RECOMMENDED DAILY INTAKE & FEEDING DURATION

ADULT DOG	IDEAL WEIGHT (kg)	STAGE			
		WEIGHT LOSS		MAINTENANCE AFTER WEIGHT LOSS	
		grams	cups	grams	cups
	2	37	4/8	41	4/8
	4	63	6/8	69	7/8
	6	85	1	93	1 + 1/8
	8	105	1 + 2/8	116	1 + 3/8
	10	125	1 + 4/8	137	1 + 5/8
	15	169	2 + 1/8	186	2 + 2/8
	20	210	2 + 5/8	230	2 + 7/8
	25	248	3	272	3 + 3/8
	30	284	3 + 4/8	312	3 + 7/8
	35	319	3 + 7/8	351	4 + 2/8
	40	352	4 + 3/8	388	4 + 6/8
	45	385	4 + 6/8	423	5 + 1/8
	50	417	5 + 1/8	458	5 + 5/8
	55	447	5 + 4/8	492	6
	60	478	5 + 7/8	525	6 + 3/8
	70	536	6 + 5/8	590	7 + 2/8
	80	593	7 + 2/8	652	8

Weight loss: Individual target weight is determined and then feeding amount established. The recommended daily intake is based on the dog's ideal weight. The starting ration may need to be adjusted during follow-up visits, based on each individual's response and the rate of weight loss. An objective of losing 1-3% bodyweight per week is generally achievable over the first 3 months, and of 0.5-2% afterwards.

Maintenance after weight loss: Energy requirements to maintain stable body weight after weight loss are lower than those of dogs that have never had obesity. Based on current scientific knowledge, our maintenance after weight loss rations are calculated by increasing the weight loss ration by about 10%. It is recommended to divide the daily intake into at least 2 meals and have fresh water available at all times. Maximum benefits on mobility support are expected after 4 to 8 weeks feeding.

ROYAL CANIN® SATIETY + MOBILITY dry is a complete dietetic feed for dogs formulated for the reduction of excessive body weight.

RECOMMENDATIONS: your advice as a veterinarian is necessary before use of this diet. Feed until target body weight is achieved and after if required to maintain target body weight. For an efficient weight loss or ideal weight maintenance, the recommended daily energy intake should not be exceeded. The diet may be continued for the dog's lifetime with vet follow-up.



RECOMMENDED FOR CASES OF

Weight management:

- Overweight/obesity: weight loss and maintenance after weight loss
- Stabilised diabetes mellitus with overweight/obesity
- Hyperlipidemia with overweight/obesity

Mobility:

- Decreased mobility
- Joints in need of support after orthopaedic surgery or trauma

NOT RECOMMENDED IN CASES OF

- Growth, gestation/lactation
- Underweight
- Conditions where high fibre content is not recommended (eg: chronic constipation/history of constipation in response to a high fibre diet)



EFFECTIVE WEIGHT MANAGEMENT

Provides safe weight loss and helps avoid weight regain. 97% of dogs lost weight in 3 months using this nutritional approach*.



BEGGING CONTROL

High level of selected fibres helps keep dogs satisfied between meals. Nutritional approach shown to help control begging in 83% of dogs during weight loss*.



LOW CALORIE

Overweight can negatively impact mobility. Low calorie content to help achieve and maintain an ideal body weight, thus supporting healthy mobility.



JOINT COMPLEX

Increased level of green-tipped mussel and high level of omega-3 fatty acids (EPA+DHA) to support mobility and help maintain healthy joints.

*Flanagan J et al (2017) Success of a weight loss plan for overweight dogs: The results of an international weight loss study. *PLoS ONE* 12(9):e0184199.

KEY VALUES

	Per 1000 kcal	Per 100 g as fed
Moisture	36.57 g	9.50 g
Protein	115 g	30 g
Fat content	40.42 g	10.50 g
Crude fibre	64.67 g	16.80 g
Ash	23.09 g	6 g
Carbohydrate (NFE)	105 g	27.20 g
Dietary fibre	108 g	28.10 g
Omega-6 fatty acids	8.01 g	2.08 g
Omega-3 fatty acids	4.50 g	1.17 g
EPA + DHA	3.08 g	0.80 g
Glucosamine + Chondroitin	770 mg	200 mg
Calcium	3.70 g	0.96 g
Phosphorus	2.81 g	0.73 g
Sodium	1.15 g	0.30 g
Chloride	3.73 g	0.97 g
Magnesium	0.31 g	0.08 g
Vitamin D	384.91 IU	100 IU
Vitamin E	277.14 IU	72 IU
Vitamin C	115.47 mg	30 mg
Metabolisable energy**		259.80 kcal

**Calculated according to NRC 2006.

COMPOSITION

Vegetable fibres, dehydrated poultry proteins, wheat gluten, manioc (tapioca), wheat, maize gluten, hydrolysed animal proteins, maize, beet pulp, animal fats, fish oil, minerals, maize flour, psyllium husks and seeds, soy oil, fructo-oligosaccharides, New Zealand green-tipped mussel powder (GLM) (0.45%), algal oil *Schizochytrium* sp. (source of EPA+DHA), glucosamine from fermentation, mango meal, hydrolysed cartilage (source of chondroitin 0.01%).

ADDITIVES (KG)***

Nutritional additives: Vitamin A: 19200 IU, Vitamin D3: 1000 IU, Iron (3b103): 30 mg, Iodine (3b201, 3b202): 3 mg, Copper (3b405, 3b406): 9 mg, Manganese (3b502, 3b504): 39 mg, Zinc (3b603, 3b605, 3b606): 132 mg, Selenium (3b801, 3b811, 3b812): 0.04 mg - Preservatives - Antioxidants.

*** Values reflect only levels added to the formula, not those naturally occurring in components of the diet.



RENAL + MOBILITY



RENAL + MOBILITY



Available in 3.5, and 9 kg bags

RECOMMENDED DAILY INTAKE & FEEDING DURATION

	DOG'S WEIGHT (kg)	BODY CONDITION					
		THIN		NORMAL		OVERWEIGHT	
		grams	cups	grams	cups	grams	cups
ADULT DOG	2	53	5/8	47	4/8	40	3/8
	4	89	1	78	7/8	68	6/8
	6	121	1 + 2/8	106	1 + 1/8	92	1
	8	150	1 + 5/8	132	1 + 3/8	114	1 + 2/8
	10	177	1 + 7/8	156	1 + 5/8	135	1 + 4/8
	15	240	2 + 5/8	211	2 + 2/8	183	2
	20	298	3 + 1/8	262	2 + 6/8	227	2 + 3/8
	25	352	3 + 6/8	310	3 + 2/8	268	2 + 7/8
	30	404	4 + 3/8	356	3 + 6/8	307	3 + 2/8
	35	454	4 + 7/8	399	4 + 2/8	345	3 + 5/8
	40	501	5 + 3/8	441	4 + 6/8	381	4 + 1/8
	45	548	5 + 7/8	482	5 + 1/8	416	4 + 4/8
	50	593	6 + 3/8	522	5 + 5/8	450	4 + 6/8
	55	637	6 + 6/8	560	6	484	5 + 1/8
	60	679	7 + 2/8	598	6 + 3/8	516	5 + 4/8
	70	763	8 + 1/8	671	7 + 1/8	580	6 + 2/8
	80	843	9	742	7 + 7/8	641	6 + 7/8

It is recommended to divide the daily intake into 2 meals and have fresh water available at all times.

Renal diets are recommended as soon as creatinine goes out of reference range (from CKD late stage 2) and/or as soon as there is proteinuria, before the disease further progresses and before inappetence develops. A food transition over 4-7 days is recommended. Maximum benefits on mobility support are expected after 6 to 8 weeks feeding.

ROYAL CANIN® RENAL + MOBILITY dry is a complete dietetic feed for dogs formulated for the support of renal function in case of chronic renal insufficiency, with restricted level of phosphorus and high quality proteins.

RECOMMENDATIONS: your advice as a veterinarian is necessary before use and before extending the period of use of this diet. For the support of renal function in case of chronic renal insufficiency, feed this diet initially up to 6 months before recheck. The diet may be continued for the dog's lifetime with vet follow-up.

RECOMMENDED FOR CASES OF

Renal disease:

- Azotaemic Chronic Kidney Disease (IRIS late stage 2 to stage 4)
- Chronic Kidney Disease with proteinuria (IRIS stage 1 to 4)

Mobility:

- Decreased mobility
- Joints in need of support after orthopaedic surgery or trauma

Other:

- Reduction of recurrence of urate or cystine urolithiasis in dogs with impaired renal function

NOT RECOMMENDED IN CASES OF

- Growth, gestation/lactation
- Pancreatitis



RENAL SUPPORT

Formulated with low phosphorus and moderate protein of high quality to support renal function, helping to improve the dog's quality of life.



HIGH ENERGY

High energy content to reduce meal volume and help compensate for a decreased appetite.



JOINT COMPLEX

With Green Lipped Mussel and omega-3 fatty acids (EPA+DHA) to support mobility and help maintain healthy joints.



AGEING SUPPORT

Highly digestible formula enriched with vitamins to support healthy ageing.

KEY VALUES

	Per 1000 kcal	Per 100 g as fed
Moisture	23.95 g	9.50 g
Protein	34.04 g	13.50 g
Fat content	44.13 g	17.50 g
Crude fibre	5.80 g	2.30 g
Ash	10.09 g	4 g
Carbohydrate (NFE)	134 g	53.20 g
Dietary fibre	20.17 g	8 g
Omega-6 fatty acids	9.61 g	3.81 g
Omega-3 fatty acids	3.05 g	1.21 g
EPA + DHA	1.66 g	0.66 g
Glucosamine + Chondroitin	276mg	110 mg
Calcium	0.98 g	0.39 g
Phosphorus	0.50 g	0.20 g
Sodium	0.88 g	0.35 g
Chloride	2.07 g	0.82 g
Magnesium	0.25 g	0.10 g
Potassium	1.51 g	0.60 g
Vitamin D	277 IU	110 IU
Vitamin E	123.55 IU	49 IU
Metabolisable energy*		396.60 kcal

*Calculated according to NRC 2006.

Essential fatty acid (linoleic acid): 3.64 %

COMPOSITION

Maize flour, rice, animal fats, wheat flour, maize gluten**, wheat, hydrolysed poultry liver, beet pulp, wheat gluten**, minerals, fish oil, vegetable fibres, soya oil, dried tomato pulp (source of lycopene), psyllium husks and seeds, sodium butyrate, fructo-oligosaccharides, New Zealand green-lipped mussel powder (GLM) (0.3%), mono- and diglycerides of palmitic and stearic acids esterified with citric acid, algal oil Schizochytrium sp. (source of EPA-DHA), glucosamine from fermentation, marigold meal, hydrolysed cartilage (source of chondroitin 0.009%).

**Protein sources: 91%.

ADDITIVES (KG)***

Nutritional additives: Vitamin A: 17 900 IU, Vitamin D3: 1100 IU, Vitamin E: 490 mg, Vitamin C: 300 mg, Iron (3b103): 45 mg, Iodine (3b201, 3b202): 4.5 mg, Copper (3b405, 3b406): 13.8 mg, Manganese (3b502, 3b504): 59 mg, Zinc (3b603, 3b605, 3b606): 153 mg, Selenium (3b801, 3b811, 3b812): 0.04 mg - Technological additives: Cinepitolite of sedimentary origin: 5 g - Preservatives - Antioxidants.

*** Values reflect only levels added to the formula, not those naturally occurring in components of the diet.





UK

WWW.ROYALCANIN.COM/UK

NUTRITIONAL HELPLINE

0800 717800

ROI

WWW.ROYALCANIN.COM/IE

NUTRITIONAL HELPLINE

1800 936074